Appendix A

The following discussion includes those projects requiring explanation beyond the summaries provided in Chapter 1 for ranking, sampling plan development, chemical testing, biological testing, or those for which the DMMP agencies used best professional judgement.

Dredging Year 2000

USACE Olympia Harbor Maintenance Dredging. The Corps of Engineers and the Port of Olympia proposed to dredge approximately 635,000 cubic yards of sediment from Olympia Harbor. There were no exceedances of 1999 DMMP screening levels for the standard list of chemicals of concern for the Olympia Harbor Maintenance project. A tiered approach was used in the analysis for Tributyltin (TBT). Composites in the berthing are (B1 and B2) were analyzed for TBT. If there were no TBT exceedances in these samples, TBT testing would not be required for the rest of the samples. Composite B1 had a porewater TBT level of 0.28 ug/L, above the screening level of 0.15 ug/L. This exceedance triggered the requirement to test all remaining DMMUs for TBT. One of these DMMU also exceeded the screening level for TBT (TBW-1). DMMU exceeding the screening level for TBT are required to undergo bioaccumulation testing in order to determine suitability for open-water disposal.

A separate sampling and analysis effort was undertaken for the bioaccumulation testing of samples B1 and TBW-1. A sampling plan addendum was approved by the agencies in July 1999. Sampling for TBT bioaccumulation analysis was completed in August 1999.

Bioaccumulation testing was performed with bivalve *Macoma nasuta* and the polychaete *Nephtys caecoides*. The two species were tested together in the same 18-liter glass aquarium. At the time of project initiation, the standard DMMP bioaccumulation protocol called for 28-day test duration. The project proponents agreed to extend the test to 45 days, based on the recommendation of the DMMP agencies. The extended test provides a better approximation of steady-state tissue concentrations for TBT.

Six replicate aquaria (five test replicates and one replicate for steady state monitoring) were run for the two test sediments, the two reference sediments and the negative control.

Tissue concentrations from the 45-day exposure were compared to the reference sediments. Initial sediment chemistry was used to adjust the observed tissue concentrations. The sediment chemistry results between the first and second rounds of TBT testing differed, and so a ratio of the two was used to adjust the bioaccumulation tissue concentrations to reflect a "worst case" analytical result. These TBT chemistry results are as follows: DMMU B1

Initial (4/99) .28(?g/L), retest (8/99) 00.14 (?g/L), ratio I/R 2.0; DMMU TBW1: Initial (4/99) 0.16 (?g/L), retest (8/99) 0.02 (?g/L), ratio I/R 8.0.

The DMMP agencies agreed to use the target tissue level developed for the East Waterway project, 3 ppm dry weight of TBT in tissue, as the value appropriate for the Olympia Harbor Navigation Project. Given the limited residue-effects data for the Olympia area, it was determined that the number calculated for Elliott Bay would be the closest approximation available for making a determination of suitability. The method of calculation and the data supporting this value is documented in the suitability determination for the East Waterway project suitability determination (1999), paragraph 18, and in the "Review of Tissue Residue Effects Data for Tributyltin, Mercury and Polychlorinated Biphenyls", prepared by EVS solutions for the Port of Seattle.

TBT concentrations in tissues from *Macoma* and *Nephtys* exposed to test sediments were significantly less than the target tissue level of 3 ppm dry weight TBT in tissue. TBT tissue concentrations were also compared to reference and significant differences were observed for both DMMU. The DMMP agencies agreed that comparing statistical difference from reference is a necessary but not a sufficient condition to determine a DMMU unsuitable for open-water disposal. Sediments from these two DMMU are suitable because all TBT tissue concentrations are significantly less than the target tissue level, TBT is undetected in most test replicates and differs from reference only when conservative assumptions are applied to non-detected values, and TBT concentrations in the retested samples were all lower than the screening level.

<u>East Waterway Stage II Project.</u> The East Waterway Project was located in the high ranked lower Duwamish Waterway. The use of best professional judgement for this project was exercised during bioaccumulation testing. The bioassay testing results were relatively routine and will not be discussed here.

Of the 49 DMMUs that had bioaccumulation trigger (BT) exceedances 25 (including S-23) passed the DMMP bioassays interpretation guidelines for openwater-unconfined disposal during Phase 1 testing. **Table A-1** (also see **Appendix C**) highlights the 25 DMMUs and chemicals exceeding BTs that were subject to bioaccumulation testing during Phase 2.

The sediment analytical results for the initial and resampled/retested sediment for the 25 DMMUs requiring bioaccumulation testing are presented in **Table A-1**. The results of these sediment analyses indicated that there was often a large disparity between the Phase 1 and Phase 2 analytical results for the COCs that exceeded the BTs. When sediment chemistry results from Phase I testing exceeded those from Phase II, the ratio of the two was used to adjust the bioaccumulation tissue concentrations to reflect a "worst case" analytical result. In the cases where the ratio was less than 1 (Phase 1 < Phase II), no adjustments were made to the tissue concentration. Conventional sediment

parameters were also reanalyzed for the 25 DMMUs and indicated the sediment characteristics were largely similar between Phase 1 and 2.

As noted above, only 25 of 49 DMMUs with one or more BT exceedances in Phase 1 testing were subjected to bioaccumulation testing. The remaining 24 DMMUs failed Tier III bioassay testing and no additional testing (e.g., bioaccumulation) was required to complete the suitability determination.

Bioaccumulation testing was performed with *Macoma nasuta*, a facultative deposit feeding/suspension feeding bivalve and *Nephtys caecoides*, a burrowing facultative deposit feeding/carnivorous polychaete. The two species were tested together in the same 8-gallon aquaria. The standard PSDDA bioaccumulation test duration is 28 days. However, to provide a better approximation of steady-state tissue concentrations for the tested chemicals (TBT, Fluoranthene, total DDT, and total PCBs), the applicant (Corps of Engineers/Port of Seattle) agreed to extend the exposure period to 45 days based on the recommendation of the DMMP agencies. The actual test was terminated at 44 days due to an increased rate of mortalities among the test species near the end of the test period.

Table A-1. DMMUs with bioaccumulation trigger exceedances.

		Sediment Concentr	ations exceeding BT	
	TBT	Fluoranthene	Total DDT	Total PCBs
	BT = 0.15	BT = 4,600	BT = 50	BT = 38
DMMUs > BT	?g/liter (porewater)	?g/kg-DW	?g/kg-DW	mg/kg -OC normDW
	Initial / Retest	Initial / Retest	Initial / Retest	Initial / Retest
	(I/R Ratio)	(I/R Ratio)	(I/R Ratio)	(I/R Ratio)
S4	0.18 / 0.11 (1.64)			
S5	0.31 MB / 0.09 (3.4)			103 / 53 (<mark>1.94</mark>)
S6	0.15 MB / 0.08 (1.9)			50 / 21 (<mark>2.4</mark>)
S7	0.19 MB / 0.09 (2.1)			
S8	0.17 M / 0.24 (0.7)			
S9				48 / 103 (<mark>0.46</mark>)
S10				42 / 329 (<mark>0.13</mark>)
S11			51 U / 47 (<mark>1.1)</mark>	127 / 42 (<mark>3.0</mark>)
S13				44 / 82 (<mark>0.54</mark>)
S14				56 / 98 (<mark>0.57</mark>)
S16			58 UJ / 61 (<mark>0.95</mark>)	77 / 44 (<mark>1.75</mark>)
S19				45 / 44 (<mark>1.02</mark>)
S21	0.15 M / 0.17 (<mark>0.9</mark>)			90 / 60 (<mark>1.5</mark>)
S23	0.28 J / 0.22 (<mark>1.3</mark>)		98 U / 43 (<mark>2.28</mark>)	212 / 81 (<mark>2.6</mark>)
S31	0.35 B / 0.51 (<mark>0.6</mark>)			
S39	0.23 M / 0.77 (0.3)			
S40	0.19 M / 1.05 (<mark>0.18</mark>)			
S41	0.23 M / 0.18 (1.3)			
S43	0.21 MB / 0.12 (<mark>1.75</mark>)			
S46	0.22 / 0.38 ((<mark>0.58</mark>)			
S47	0.83 / 4.0 (0.21)			
S49	0.25 MB / 0.24 (1.04)			38 / 90 (<mark>0.42</mark>)
S50	0.19 B / 0.12 (<mark>1.6</mark>)	6,400 / 800 (<mark>8.0</mark>)		88 / 41 (<mark>2.15</mark>)
S52	0.20 M / 0.17 (<mark>1.2</mark>)			
S57	0.92 MB / 0.47 (<mark>1.96</mark>)	·		

Legend: DW = dry weight; OC = organic carbon normalized value; M = estimated value; B = possible blank contamination; J = estimated value; U = Undetected at reported concentration; UJ = analyte not detected above the reported sample quantitation limit; Shaded cells denote DMMU's failing bioaccumulation test interpretation guidelines, which are discussed in text; I/R Ratio = highlighted yellow value: ratio of initial/retested for tested analyte.

Five replicate 8-gallon aquaria were run for the negative control, for each of the 3 reference sediments, and for each of the 25 tested DMMUs. In addition to the routine water quality metrics (temperature, salinity, dissolved oxygen, pH) that were monitored during the exposure period, the DMMP agencies recommended and the applicant agreed to collect an additional metric, wetweight growth, during the exposure period to further assess the general health and well-being of the test animals. The results of growth and survival measurements taken for each species during the exposure period suggested

that for *Macoma nasuta* there was no apparent relationship between mean growth and survival during the exposure period. The results for *Nephtys caecoides* indicated there was a statistically significant (p<0.01) negative effect on survival with a reduction in mean growth during the exposure period.

Tissue concentrations of chemicals-of-concern from the 44-day exposures were compared statistically to the appropriate reference sediment, based on grain size similarity comparisons. As noted above, the calculated ratios of Phase 1 (initial)/Phase II (retest) sediment chemistry were used to adjust the observed tissue concentrations. Statistical comparisons of test DMMUs and reference tissue concentrations for the final interpretation "worst case" analyses were based on the adjusted tissue concentrations. The summary tissue chemistry interpretation for each of the measured chemicals is provided in **Table A-2** for each of the 25 DMMUs tested.

The DMMP agencies agreed that comparing statistical differences from reference is a necessary, but not sufficient condition to determine a DMMU unsuitable for open-water disposal. For those DMMUs that were statistically greater than reference, a more in depth evaluation was required to determine the significance of the bioaccumulation that had occurred. This evaluation focused on **a)** Food and Drug Administration (FDA) Action Levels for Poisonous and Deleterious Substances in Fish and Shellfish for Human Food; **b)** PSDDA target tissue concentration values for chemicals of concern to human health, and **c)** ecological residue-effects data from the literature.

a) The FDA guidelines for the chemicals of concern addressed by East Waterway Stage II bioaccumulation testing are as follows:

Tributyltin (TBT): No guideline Fluoranthene: No quideline

DDT + DDE: 5.0 ppm wet weight (ww)

PCBs: 2.0 ppm ww

b) A risk-based approach was adopted by the PSDDA program in 1988 to set target tissue levels (TTL) for human health. The TTL calculated for **fluoranthene** based on risk to humans consuming seafood is **8,400 ppm wet weight**.

As part of a suitability determination for the Port of Seattle T-18 dredging project (March 17, 1997 SDM), the PSDDA agencies reevaluated the human health-based TTLs for PCBs, total DDT, mercury, and TBT. In recalculating these TTLs, the PSDDA agencies used updated cancer slope factors and reference doses, as well as estimates of fish home range. The TTL developed for total DDT is 44 ppm wet weight.

The DMMP agencies recently undertook a re-evaluation of the PCB TTL for human health. Recalculation of the PCB TTL for the Elliott Bay disposal site included using an updated cancer slope factor, recent fish consumption data, and consideration of PCB biomagnification due to trophic transfer. Based on this analysis, an interim TTL for total PCBs (Aroclor) of 0.75 ppm wet weight has been used to interprete bioaccumulation data from the East Waterway Phase II Project.

A recent effort by the Port of Seattle (May 1999)¹ involved compilation of the residue-effect literature for TBT. It was prepared for the Port of Seattle by EVS Solutions for submittal to the U.S. Environmental Protection Agency for the Harbor Island Superfund Site, Waterway Sediment Operable Unit. Using residue-effects data from this and other studies, EPA Superfund developed a tissue trigger level of 3 ppm dry weight of TBT in tissue (0.6 ppm wet weight) that was used to evaluate bioaccumulation data from the West Waterway OU (for more information see Appendix D of the May 1999 EVS report). This tissue concentration is protective for growth and reproduction endpoints in polychaetes, crustaceans, bivalves, and most gastropods. However, it might not protect the most sensitive species of meso- and neogastropods against imposex-related sterility. Considering that meso- and neogastropods are rare in Elliott Bay (Appendix D in EVS, 1999), the DMMP agencies have decided to use the West Waterway TBT trigger level (3 ppm dry weight) on an interim basis to interpret bioaccumulation relative to disposal at the Elliott Bay site.

To summarize, the DMMP agencies used the following TTLs to interpret the bioaccumulation test data for the East Waterway Stage II Project:

TBT: 3.0 ppm dry weight (dw) as TBT

Fluoranthene: 8,400 ppm ww DDT + DDE: 3.0 ppm ww PCB: 0.75 ppm ww

The agencies used best professional judgement in developing these interpretation guidelines to meet PSDDA disposal site management objectives; achievement of other sediment management objectives will require additional evaluation. These guidelines are subject to change for future PSDDA/DMMP projects as additional bioaccumulation data become available.

Each DMMU was compared to these interpretation guidelines using a one-tailed one-sample t-test (see **Table A-2**). An alpha level (the probability of making a Type I error, rejecting the null hypothesis of no difference between test and

_

¹ For TBT, the DMMP agencies relied upon Appendix D of a May 1999 report entitled: "Review of Tissue Residue Effects Data for Tributyltin, Mercury, and Polychlorinated Biphenyls". Prepared by EVS Solutions for the Port of Seattle.

reference responses when, in fact, they are not different) of 0.1 was selected for these statistical comparisons by the DMMP agencies to reflect the higher within sample variability, and to increase the power of the test to discriminate between reference and test responses. DMMU S23 exceeded the interim PCB target tissue level (TTL) and thus failed the bioaccumulation test. Because S23 (ML Rule exceedance) failed the bioaccumulation test for PCBs, further testing under a Tier IV evaluation is unnecessary to make a suitability determination. Two additional DMMUs, S11 and S16 statistically exceeded the interim PCB TTL, and also failed the bioaccumulation test. DMMU S31 exceeded the interim TBT TTL and therefore failed the bioaccumulation test. No other DMMUs statistically exceeded the bioaccumulation interpretation guidelines. In summary, of the 25 DMMUs tested representing 95,340 cubic yards, 4 DMMUs failed the bioaccumulation test representing a total volume of 15,680 cubic yards.

Chemical/biological testing conducted for a portion of the East Waterway Stage II Project surfaced subsurface contamination issues that will require further examination of the subsurface sediment quality (e.g., Z-sample collection and analysis) of the proposed new-sediment surface following dredging to verify compliance with Washington State's antidegradation statute. These analyses have not yet been conducted and must be completed before dredging of those identified DMMUs can be conducted.

Table A-2. Worst Case Bioaccumulation Interpretation Summary (Adjusted values)

							MMU S	4									MMU S	55								D	OMMU S	6				
				Ma	coma na	suta			Neph	tys caed	coides			Ma	coma n	asuta			Neph	tys caed	oides			Ma	coma na	suta			Neph	tys caec	oides	
TBT ion (as TBT) up	ug/kg-dw ug/kg-ww	Guideline 3,000 8,400	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	98 DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	65 DMMU tissue (Initial)	Sp DMMU tissue (adjusted) ය	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline
	ıg/kg-ww	3,000										-	334	444	0.00			007	200	45.0			400	544	0.00			400	529	45.0		
Total PCBs uç	ug/kg-ww	750				Г	MMU S	7					334	441	9.88		yes MMU S	607 8	802	15.9	yes	yes	189	511	9.88	yes	yes OMMU S	196	529	15.9	yes	yes
				Ma	coma na		, minio 0	İ	Nenh	tys caed	roides			Ma	coma n			Ĭ	Nenh	tys caed	oides			Mai	coma na				Nenh	tys caec	nides	
TBT ion (as TBT) ug Fluoranthene ug Total DDT ug	ug/kg-dw ug/kg-ww ug/kg-ww	Guideline 3,000 8,400 3,000	PMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	9 Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	E Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	US DMMU tissue (Initial)	DMMU issue (adjusted)	99 Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (SB-Mac Control)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (SB-Mac Control)	Statistically different from reference	statistically below guideline
Total PCBs ug	ıg/kg-ww	750																					340	340	17.2	yes	yes	615	615	17.3	yes	yes
							MMU S1	10									MMU S	11									MMU S	13				
				ма	coma na	isuta		—	Neph	tys caed	oides	1	 	Ма	coma n	asuta		-	Neph	tys caed	olaes	1		Ма	coma na	isuta		—	Neph	tys caec	coldes	
CHEMICAL NAME	Units	Guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (SB-Mac Control)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (SB-Mac Control)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (SB-Mac Control)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (SB-Mac Control)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (SB-Mac Control)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (SB-Mac Control)	Statistically different from reference	statistically below guideline
	ug/kg-dw	3,000					,,																									
	ıg/kg-ww	8,400												•																		
	ug/kg-ww	3,000	200	200	17.0	1/00	V00	750	750	17.0	1105	VOC	16.5 490	18.0 1.532	0.44	yes	yes	21.5	23.4 2.036	0.76	yes	yes	217	217	17.0	1/00	WOO	471	474	17.0	V00	V00
Total PCBs uç	ıg/kg-ww	750	398	398	17.2	yes	yes	750	750	17.3	yes	yes	490	1,532	17.2	yes	no	651	2,036	17.3	yes	no	217	217	17.2	yes	yes	4/1	471	17.3	yes	yes

Note: (1) All tissue concentrations for Fluoranthene, Total DDT and Total PCBs were converted to wet weight to facilitate guideline comparisons. All TBT tissue concentrations are on a dry weight basis. (2) Adjustments to tissue concentrations based on initial sediment versus retested sediment concentration ratios (see Appendix 5). Concentration ratios greater than 1 were adjusted.

Concentration ratios less than 1 were not adjusted.

Table A-2. Worst Case Bioaccumulation Interpretation Summary (Adjusted values)

				DI	MMU S	14								D	MMU S1	6								D	MMU S1	9				
		Ma	coma na	asuta			Nephi	tys cae	coides			Ma	coma na	suta			Nephi	tys caec	oides			Ma	coma na	asuta			Neph	tys caec	oides	
CHEMICAL NAME Units Guideline	DMMU tissue (Initial)	OMMU tissue (adjusted)	eference (SB-Mac Control)	Statistically different from reference	tatistically below guideline	JMMU tissue (Initial)	MMU tissue (adjusted)	kieference (SB-Mac Control)	Statistically different from reference	tatistically below guideline	DMMU tissue (Initial)	OMMU tissue (adjusted)	seference (SB-Mac Control)	Statistically different from reference	tatistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	seference (SB-Mac Control)	Statistically different from reference	tatistically below guideline	OMMU tissue (Initial)	MMU tissue (adjusted)	(cference (CR-23 Mod)	Statistically different from reference	tatistically below guideline	oMMU tissue (Initial)	DMMU tissue (adjusted)	Kaference (CR-23 Mod)	Statistically different from reference	statistically below guideline
TBT ion (as TBT) ug/kg-dw 3,000			-	0,	0)			ш.	0)	0)		_	- 12	0)	63			ш.	0)	0)				0,	0)			-	0)	0)
Fluoranthene ug/kg-ww 8,400																														
Total DDT ug/kg-ww 3,000				1							18.4	18.4	0.44	yes	yes	36.2	36.2	0.76	yes	yes									├	
Total PCBs ug/kg-ww 750	385	385	17.2		yes MMU S	800	800	17.3	yes	yes	507	994	17.2	yes	no MMU S2	912	1,787	17.3	yes	no	233	238	9.88	yes	yes MMU S3	437	446	15.9	yes	yes
		Mo	icoma na		IVIIVIU 3	<u> </u>	Nonhi	tys cae	ooidoo			Mo	coma na		IVIIVIU 52	ა	Nonhi	tys caed	oidoo			Mo	coma na		WINO 53) I	Nonh	tys caec	noidee	
CHEMICAL NAME Units Guideline TBT ion (as TBT) ug/kg-dw 3,000 Fluoranthene ug/kg-ww 8,400 Total DDT ug/kg-ww 3,000	8. EDMMU tissue (Initial)	DMMU tissue (adjusted)	22. Reference (SB-Mac Control)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (SB-Mac Control)	Statistically different from reference	statistically below guideline	4.02 DMMU tissue (Initial)	9. 0. 22. (adjusted.) 2. 6. 6. (adjusted.) 6. 6. 6. (adjusted.) 6. 6. (adjusted.)	15.0 16.0 17.0 18.0 18.0 18.0 18.0 18.0 18.0 18.0 18	Statistically different from reference	statistically below guideline	127 54.2	DMMU tissue (adjusted)	0.76 Beference (SB-Mac Control)	Statistically different from reference	se statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	8.02 Reference (CR-23 W)	Statistically different from reference	Statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	6 Reference (CR-23 W)	Statistically different from reference	statistically below guideline
Total PCBs ug/kg-ww 750	315	337	17.2		yes	577	618	17.3	yes	yes	780	1,561	17.2	yes	no	1,575	3,150	17.3	yes	no									ш	
					MMU S	39									MMU S4	0									MMU S4	11				
CHEMICAL NAME Units Guideline TBT ion (as TBT) ua/kq-dw 3,000	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod) 8u au	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 W) eu	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 W)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod) 33.	Statistically different from reference	DMMU tissue (adjusted)	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod) sopono	Statistically different from reference	statistically below guideline
TBT ion (as TBT) ug/kg-dw 3,000 Fluoranthene ug/kg-ww 8,400	134	134	33.4	yes	yes	40.7	45.7	65.0	no	yes	204	284	20.3	yes	yes	57.8	57.8	72.9	no	yes	222	284	33.4	yes	yes	44.2	56.6	65.0	no	yes
Total DDT ug/kg-ww 3,000																														
Total PCBs ug/kg-ww 750																														

Table A-2. Worst Case Bioaccumulation Interpretation Summary (Adjusted values)

						D	MMU S	43								D	MMU S	16								D	MMU S4	1 7				
				Ma	coma na	asuta			Neph	tys caec	oides			Mad	coma na	asuta			Neph	itys caed	coides			Ma	coma na	asuta			Neph	tys caec	oides	
CHEMICAL NAME		Guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline
TBT ion (as TBT)	ug/kg-dv		71.6	91.6	33.4	no	yes	73.0	93.4	65.0	no	yes	1,220	1,220	33.4	yes	yes	294	294	65.0	yes	yes	1,780	1,780	33.4	yes	yes	352	352	65.0	yes	yes
Fluoranthene Total DDT	ug/kg-ww			1																												
Total PCBs	ug/kg-wv																															
		•				D	MMU S	49								D	MMU S	50								D	MMU S	52				
				Ма	coma na	asuta			Neph	tys caec	oides			Мас	coma na	asuta			Neph	itys caed	coides			Ma	coma na	asuta			Neph	tys caec	oides	
CHEMICAL NAME TBT ion (as TBT) Fluoranthene Total DDT	ug/kg-dw ug/kg-ww ug/kg-ww	8,400 3,000	DMMU tissue (Initial)		Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	88 51 8 87 8 87 8 88 51 8 87 8 88 51	DMMU tissue (adjusted)	20.5 8. Reference (CR-23 Mod)	Statistically different from reference	s statistically below guideline	99 DMMU tissue (Initial)	DMMU tissue (adjusted)		Statistically different from reference	statistically below guideline	Ф DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline
Total PCBs	ug/kg-wv	750	246	246	9.88		yes	590	590	15.9	yes	yes	246	523	9.88	yes	yes	364	775	15.9	yes	yes										
				14-	coma na		MMU S	5/	Mach	tys caec	-:		ł																			
CHEMICAL NAME TBT ion (as TBT)	Units ug/kg-dw ug/kg-w		DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference (CR-23 Mod)	Statistically different from reference	statistically below guideline																				
Total DDT	ug/kg-wv			1																												
Total PCBs	ug/kg-wv												1			Target	Tissue	Guidelii	ne excee	ded												

<u>U.S. Navy Puget Sound Naval Shipyard (PSNS) Pier Construction and Dredging Project</u>. The PSNS project was located in the high ranked Sinclair Inlet within a Superfund cleanup site footprint. The following biological testing summary covers the use of best professional judgement exercised by the DMMP agencies during sediment testing for this project.

During Phase I testing, the Navy elected to conduct concurrent bioassay testing on all eighty-three DMMUs (Phase I + Phase II DMMUs totaled 91 due to addition of three additional DMMUs and two subsurface DMMUs, D4 and D6, reanalyzed as 7 uncomposited DMMUs) because of timing considerations and concerns about exceeding bioassay holding times. The results of the Phase I and II bioassay results are summarized below.

Standard bioassay testing was conducted on all eighty-three Phase I DMMUs within the 56 day biological holding time. Ten DMMUs sampled during Phase II (e.g., 3 new DMMUs and 7 retested subsurface DMMUs) were subject to concurrent bioassay testing with the standard bioassay testing suite (amphipod, bivalve larval, and *Neanthes* growth bioassays). The three surface DMMUs from the Turning basin/Inner Channel (\$101, \$102, \$103) used *Ampelisca abdita* in lieu of *Eohaustorius estuarius* due to high clay contents (27 - 36.4%). The seven retested Pier subarea subsurface DMMUs were tested using both amphipod species. The DMMP agencies specified retesting with *Eohaustorius estuarius*, and the Navy also elected to run Ampelisca abdita, when sediment conventional results indicated that two of the seven uncomposited DMMUs had clay contents greater than 15 percent. Sixty-two DMMUs were subject to amphipod bioassay retesting with *Ampelisca abdita*, whereas a subset of 10 DMMUs were also retested with *Eohaustorius estuarius* bioassay to reconfirm the clay sensitivity relationship noted during Phase I testing. Appendix B summarizes the solid phase bioassay Quality Control (QC) performance guidelines as well as the solid phase bioassay interpretative guidelines for nondispersive sites, which were used to evaluate the bioassay data discussed below. **Appendix C** (pages C8-C14) summarizes the Phase I/Phase II bioassay toxicity testing outcomes for the tested DMMUs. Three reference samples were collected from Carr Inlet to block for grain size effects. In general, all negative control and reference sediments met the DMMP performance limits for each of the four bioassay tests (e.g., 2 amphipod species, bivalve larval, and *Neanthes*) to assess toxicity. Summary results (Phase 1 + Phase II) for each bioassay test are depicted in Table 1 for each of the three subareas characterized relative to the DMMP nondispersive interpretative guidelines. These bioassay results are discussed below for each of the bioassay tests.

As noted in **Table A-3**, the amphipod bioassay results tested during Phase I with *Eohaustorius estuarius* showed significant toxicity being expressed among the 83 DMMU. This was later reconfirmed during Phase II retesting testing with a 10 sample subset of the retested DMMUs. An examination of potential

nontreatment effects suggested that clay content may have been a contributing factor to the observed toxicity during Phase I, and this hypothesis was reconfirmed during Phase II testing for a small subset of samples retested with *Eohaustorius*. The possibility of high clay contents affecting *Eohaustorius* survival was sufficient for the DMMP agencies to authorize a retest of 62 DMMUs, in which clay contents exceeded 15 percent (See **Appendix C**, pages C8-C14). Subsequently, both the Phase I and II results for *Eohaustorius* were used for decision-making in the suitability determination, when the clay contents in tested sediment were less than 15 percent. Fifteen percent clay represents the upper end in the range of the reference sediments tested (11.7 -12.3 %), where relatively low toxicity was observed. Correspondingly, results for *Eohaustorius* were not used for decision-making when the clay contents were greater than 15%. Other factors, such as test animal acclimation (salinity, temperature, etc.) were evaluated during the Phase II retest and found not to be a significant factor contributing to the observed toxicity.

The DMMP agencies also evaluated alternative hypotheses for the toxicity by conducting chemical testing utilizing Atomic Emission Detector (AED) screening analyses on six surface and two subsurface samples from the Piers D, B, and 3 subareas. These analyses identified and quantified two potentially toxic components, total phosphate compounds and tricresyl phosphate. These analyses also documented the presence of a petroleum product that is lighter than motor oil and heavier than diesel fuel. These data suggest that other potentially toxic substances in the pier subarea sediments may also have contributed to the toxicity observed during Phase I testing.

As noted in **Table A-3** and **Appendix C** (pages C8-C14), the amphipod bioassay retesting of 62 DMMUs, and 7 uncomposited subsurface DMMUs (D4 and D6), and of 3 new DMMUs from the Turning Basin/Inner Channel with *Ampelisca abdita*, showed no toxity with all 72 DMMUs tested passing the nondispersive site disposal guidelines.

The results of the larval bivalve test (**Appendix C** and **Table A-3**) also showed significant toxicity being expressed although not to the extent shown in the amphipod bioassay. A total of 58 Phase I DMMUs had two-hit and 4 exhibited one-hit toxicity responses, with 21 DMMUs exhibiting no-hit responses according to the nondispersive interpretive disposal guidelines. The results of the Phase II testing, where 10 DMMUs were subject to bivalve larval bioassay testing, appeared to mimic the Phase I results, with 7 two-hit responses and 3 no-hit responses. An examination of potential clay effects on the larval bivalve toxicity after removing the Pier data did show a significant but weak correlation (r = 0.46, p < 0.001, n = 62) for the Turning Basin/Inner Channel, where fewer chemical guideline exceedances were noted. No retesting of the Sediment Larval bioassay was authorized by the DMMP agencies and the Phase I and II results were used in the DMMP non-dispersive site regulatory decision.

Overall interpretation of the bioassay responses combining the Phase I and II responses (See **Appendix C** and **Table A-3**) indicates that virtually all the material from the Turning Basin/Inner Channel passed the unconfined-openwater disposal guidelines (52 of 53 passing), whereas 18 out of 26 Pier surface DMMUs and 9 of 12 subsurface DMMUs passed the DMMP guidelines. Five of the Pier Surface DMMUs (S51, S25, S55, S61, S70) passed the bioassay testing guidelines, and were subject to required bioaccumulation testing because of bioaccumulation trigger exceedances. These DMMUs are discussed below.

Table A-3. Phase I and II (parenthesis) Bioassay interpretation summary for each PSNS subarea.

Amphipod Bioassay: (Eohaustorius estuarius)	Two-Hit Phase I (Phase II)	One-Hit Phase I (Phase II)	No-Hit Phase I (Phase II)	Total: Phase I (Phase II)		
Turning Basin (surface)	11 (1)	39 (4)	0 (0)	50 (5)		
Clay < 15 %:	0	0	0 (0)	0		
(see footnote 2 below) ²	v		V	v		
Piers D, B, 3 (surface)	5 (2)	17 (0)	4(2)	26 (4)		
Clay < 15 %:	4	5	4	13		
Piers D, B, 3 (subsurface)	2(1)	4(1)	1 (6)	7 (8)		
Clay < 15 %:	1	1	6	8		
Subtotal:	18 (4)	60 (5)	5 (8)	83 (17)		
Clay < 15%:	5	6	10	21		
Amphipod Bioassay:	Two-Hit	One-Hit	No-Hit	Total:		
(Ampelisca abdita)	Phase II only	Phase II only	Phase II only	Phase II only		
Turning Basin (surface)	0	0	49	49		
Piers D, B, 3 (surface)	0	0	15	15		
Piers D, B, 3 (subsurface)	0	0	8	8		
Subtotal:	0	0	72	72		
Bivalve Larval Bioassay: (Mytilus galloprovincialis)	Two-Hit Phase I (Phase II)	One-Hit Phase I (Phase II)	No-Hit Phase I (Phase II)	Total: Phase I (Phase II)		
Turning Basin (surface)	33 (3)	0(0)	17 (0)	50 (3)		
Piers D, B, 3 (surface)	19 (0)	3 (0)	4(0)	26 (0)		
Piers D, B, 3 (subsurface)	6 (4)	1 (0)	0(3)	7 (7)		
Subtotal:	58 (7)	4 (0)	21 (3)	83 (10)		
Neanthes Bioassay:	Two-Hit	One-Hit	No-Hit	Total:		
(Neanthes arenaceodentata)	Phase I (Phase II)	Phase I (Phase II)	Phase I (Phase II)	Phase I (Phase II)		
Turning Basin (surface)	2(0)	0 (0)	48 (3)	50 (3)		
Piers D, B, 3 (surface)	2 (0)	0 (0)	24 (0)	26 (0)		
Piers D, B, 3 (subsurface)	1 (0)	0 (0)	6 (7)	7 (7)		
Subtotal:	5 (0)	0 (0)	78 (10)	83 (10)		
DMMP Bioassay						
Determination:	Number of Sui	itable DMMUs	Number of Unsuitable DMMUs			
(Phase I + Phase II)						
Turning Basin (surface)	5	52	1			
			8			
Piers D, B, 3 (surface)		8				
	(8 9 9	,	8 3 2		

Bolded value denotes the number of DMMUs observed during Phase I testing with clay contents < 15 %, for which *Eohaustorius* bioassay results were used to make a final suitability determination. The remaining *Eohaustorius* results with clay contents > 15 % were not used in the final suitability determination.

As noted previously, a number of DMMUs passing bioassay interpretive guidelines, also exceeded bioaccumulation triggers. These are highlighted in **Table A-4** below. Additionally, two DMMUs (S77, S78) with mercury BT/ML exceedances do not need to be considered further because they are unsuitable based on bioassay testing results. Another DMMU (\$72) had a Fluoranthene BT exceedance, but also failed bioassy testing guidelines, and no further testing is necessary. The DMMP agencies deliberated on S51, which passed Phase I bioassays, but also exceeded the ML rule, and decided that standard bioaccumulation testing with a 45-day exposure would provide sufficient data in a Tier IV evaluation to enable a DMMP suitability determination to be completed. The Navy elected to proceed with bioaccumulation testing on S51, and the results of the testing of this DMMU is described below. The Navy decided not to archive bioaccumulation samples for S25, S55, S61, S70 during the Phase II resampling effort. After reviewing the Phase II bioassay retesting results, \$25, \$55, \$61, and \$70 passed the bioassay interpretative guidelines (see **Appendix C**), but the Navy decided not to pursue bioaccumulation testing of these four DMMUs because of schedule and timing considerations. Therefore, these four DMMUs are considered unsuitable for unconfined-open-water disposal based on DMMP BPJ.

Table A-4. DMMUs passing bioassays with sediment bioaccumulation trigger exceedances.

Chemical	S51	S25	S55	S61	S70
Mercury (BT = 1.5 ppm)	1.75				
Silver (BT = 6.1 ppm)	<mark>6.5</mark>				
TBT (BT = 0.15 ppb-porewater)		0.15		0.17	
DDT (BT =50 ppb)	<mark>748</mark>		96.9		
Pentachlorophenol (BT = 504 ppm)	620				
Bis(2-ethylhexyl)phthalate (BT =13,870 ppb)					31,000

As noted in paragraph above only one of the eight DMMUs with bioaccumulation trigger exceedances (S51) was subjected to bioaccumulation testing. The remaining seven DMMUs either failed Tier III bioassay testing (S72, S77, S78), or failed to conduct bioaccumulation testing as required for unconfined openwater disposal (UCOWD) consideration (S25, S55, S61, S70). Therefore, these

seven DMMUs are unsuitable for UCOWD. The sediment analytical results of the resampled DMMU S51 are depicted in **Table A-5**.

Table A-5. Comparative Initial/Resampled Chemical Sediment Concentrations evaluated during 45 day Bioaccumualtion Test of DMMU S51

CHEMICAL NAME	Units	Sediment Initial	Sediment Resample/Retest	Initial/Retest ratio:
Mercury (Hg)	mg/kg-dw	1.75	2.95	0.59
Silver (Ag)	mg/kg-dw	6.5	4	<mark>1.63</mark>
Pentachlorophenol (PCP)	ug/kg-dw	620	510	1.22
Total DDT	ug/kg-dw	748	15	<mark>49.9</mark>

The bioaccumulation testing was performed with *Macoma nasuta*, a facultative deposit feeding/suspension feeding bivalve and *Nephtys caecoides*, a burrowing facultative deposit feeding/carnivorous polychaete. The two species were tested together in the same 10-gallon aquaria. However, to provide a better approximation of steady-state tissue concentrations for the tested chemicals (mercury. silver, DDT, pentachlorophenol), the applicant agreed to extend the exposure period from 28 to 45 days based on the recommendation of the DMMP agencies.

Five replicate 10-gallon aquaria were run for the negative control, reference sediment, and for the tested DMMU S51.

Tissue concentrations of chemicals-of-concern measured during the 45-day exposures were compared statistically to the appropriate reference sediment, based on grain size similarity comparisons. The calculated ratios of Phase 1 (initial)/Phase II (retest) sediment chemistry were used to adjust the observed tissue concentrations. When sediment chemistry results from Phase I testing exceed those from Phase II, the ratio of the two are used to adjust the bioaccumulation tissue concentrations to reflect a "worst case" analytical result (Phase I / Phase II > 1). When the ratio is less than 1 (Phase 1 / Phase II < 1), no adjustments are made to the tissue concentration. Statistical comparisons of the test tissue (DMMU S51) and reference tissue concentrations for the final interpretation "worst case" analyses were based on the adjusted tissue concentrations. The summary tissue chemistry interpretation for each of the measured chemicals is provided in **Table A-6** for the four chemicals tested in DMMU S51.

The DMMP agencies agreed that comparing statistical differences from reference is a necessary, but not sufficient condition to determine a DMMU unsuitable for open-water disposal. For each chemical measured within

DMMU-S51 that were statistically greater than reference, a more in depth evaluation is required to determine the significance of the bioaccumulation that had occurred. This evaluation focused on **a)** Food and Drug Administration (FDA) Action Levels for Poisonous and Deleterious Substances in Fish and Shellfish for Human Food; **b)** PSDDA target tissue concentration values for chemicals of concern to human health, and **c)** ecological residue-effects data from the literature.

a) The FDA guidelines for the chemicals of concern addressed by the bioaccumulation testing are as follows:

mercury (methyl mercury): 1.0 ppm wet weight (ww)

silver: no guideline

DDT + DDE: 5.0 ppm wet weight (ww)

Pentachlorophenol: no guideline

- b) A risk-based approach was adopted by the PSDDA program in 1988 to set target tissue levels (TTL) for human health. The TTL calculated for **silver** based on risk to humans consuming seafood is **200 ppm wet weight**, and is **900 ppm wet weight** for **Pentachlorophenol**. As part of a suitability determination for the Port of Seattle T-18 dredging project (March 17, 1997 SDM), the DMMP agencies re-evaluated the human health-based Target Tissue Levels (TTLs) for PCBs, total DDT, mercury, and TBT. In recalculating these TTLs, the PSDDA agencies used updated cancer slope factors and reference doses, as well as estimates of fish home range. The TTL developed for **mercury is 450 ppm wet weight**, and for **total DDT is 44 ppm wet weight**.
- c) The DMMP agencies conducted a literature review of ecological effects, relative to tissue concentrations (1997 Port of Seattle T-18 (East Waterwau Stage I) Suitability Determination). The agencies reviewed the literature data in the context of DMMP site management objectives:

"The biological testing guidelines for Site Condition II, which allow for minor significant effects in the laboratory tests, suggest that some biological effects may be expected at the disposal site. The severity and extent of biological effects are not expected to be great because the majority of the species found at the preferred disposal sites are not known to be acutely sensitive to chemicals of concern. Effects associated with Site Condition II will include sublethal effects and, potentially, an increase in the mortality of the more sensitive but less abundant crustacean species. Cumulative effects are expected to consist of a reduction in population and community biomass and an increase in the tissue concentration levels of chemicals of concern."

 Table A-6. Bioaccumulation Testing Summary for DMMU S51.

							DMMU S51					
					<i>Aacoma</i>					Nephty caecoia		
					nasuta							
CHEMICAL NAME	Units	TTL Guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Carr Inlet Reference	Statistically different from reference	statistically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Carr Inlet Reference	Statistically different from reference	statistically below guideline
Mercury (Hg)	mg/kg- ww	1	0.51	0.51	0.12	yes	yes	0.015	0.015	0.038	no	yes
Silver (Ag)	mg/kg- ww	200	0.58	0.95	0.21	yes	yes	0.055	0.089	0.055	no	yes
Pentachlorophenol (PCP)	ug/kg- ww	900	57.0	69.3	51.6	no	yes	41.3	50.2	40.1	no	yes
Total DDT	ug/kg- ww	3,000	48.7	2,429	4.0	yes	yes	58.7	2,927	0.54	yes	yes

- d) It was clear from the literature review, that for mercury, human health concerns occurred at lower tissue concentrations than ecological effects. Thus, the DMMP agencie used the FDA guideline (1.0 ppm ww) as the mercury TTL. Conversely, for DDT, ecological effects are expected to occur at a lower concentration than human health effects. A literature review conducted as part of the T-18 dredging project identified a concentration range of 3-5 ppm ww in gonads or liver for croakers and cutthroat trout associated with induction of sterility and other reproductive effects³. Therefore, the DMMP agencies selected 3.0 ppm ww for the total DDT TTL to comply with Site Condition II.
- e) To summarize, the DMMP agencies used the following TTLs to interpret the bioaccumulation test data for DMMU-S51:

Mercury (methyl mercury): 1.0 ppm ww Silver: 200 ppm ww DDT + DDE: 3.0 ppm ww Pentachlorophenol: 900 ppm ww

The agencies used best professional judgement in developing these interpretation guidelines to meet DMMP disposal site management objectives. These guidelines are subject to change for future DMMP projects as additional bioaccumulation guidance become available.

Each DMMU was compared to these interpretation guidelines using a one-tailed one-sample t-test. An alpha level (the probability of making a Type I error, rejecting the null hypothesis of no difference between test and reference responses when, in fact, they are not different) of 0.1 was selected for these statistical comparisons by the DMMP agencies to reflect the higher within sample variability, and to increase the power of the test to discriminate between reference and test responses. All four chemicals measured in S51 *Macoma* and *Nephtys* tissue were judged to be statistically below the chemical quideline using the adjusted values.

However, the results of the sediment reanalysis of S51 indicated that there was a large disparity between the Phase 1 and Phase 2 analytical results for the COCs that exceeded the BTs, especially for DDT, which was measured at 2 percent of the Phase I result (**Table A-5**). The sediment analysis results

Childress, R., Texas Parks and Wildlife Department, Levels of concentration and incidence of various pesticide residues in Texas. (unpublished report, 1971).

³ Allison, D.B., B.J. Kallman, O.B. Cope, and C.C. Van Valin. 1964. Some chronic effects of DDT on cutthroat trout. Washington, DC.: U.S. Fish and Wildlife Service, Bureau of Sport Fish. 30 pp.

indicated that silver, PCP and DDT all had ratios greater than 1, and the tissue concentrations were adjusted accordingly (Table A-6), whereas mercury was less than 1 and no adjustment was made. Because of the 50-fold discrepancy in the total DDT sediment concentration between the retested (15 ppb) and the initial sediment (748 ppb), the DMMP agencies had serious concerns about the validity of the DDT concentrations measured in the tissues of *Macoma* and Nephtys. The agencies were concerned that the DDT bioaccumulation observed in the retested sediments was not a realistic evaluation of bioavailable DDT associated with this DMMU. The discrepancy in sediment DDT was only brought to the attention of the DMMP agencies after the bioaccumulation test had been completed. The DMMP agencies deliberated and concurred that the data for DDT was insufficient for regulatory decision-making and that the test would have to be repeated using a higher DDT exposure concentration closer to the Initial Phase I concentration to be valid for decision-making. The Navy when informed of this decided not to resample and retest S51. Therefore, DMMU-S51 was considered unsuitable for UCOWD using BPJ.

USACE Duwamish Maintenance Dredging. The Corps routinely dredges only the turning basin and upper portion of the Duwamish River Navigation Channel, where low-ranked sediment deposits from the upper reaches of the river. The 2000 characterization was for the lower part of the navigation channel, where sedimentation rates are much lower but contamination issues are greater. All material sampled and tested as part of this characterization was high ranked, and though contamination was found and a portion of the proposed dredged material found unsuitable for open water disposal, for the most part the testing was routine.

The issue of note for this characterization was that a mapping error caused some samples to be taken outside the dredging prism. The error did not occur at the time of sampling, but at the time of Sampling and Analysis Plan (SAP) preparation. With no recent bathymetry maps available, the contractor used cross sections generated from the latest Corps survey to propose sampling locations. Unfortunately, those cross sections were interpreted "backwards," as if one were looking downstream, rather than upstream. In places were shoaling was evident on only one side of the navigation channel, this error resulted in a SAP with samples placed on the opposite side of the channel from the actual shoaling.

The SAP was approved by the DMMP agencies, sampling and analysis took place, and a suitability determination (3 February 2000) was signed. It was only when actual planning for the dredging took place that the error was discovered. The problem affects nine out of 20 DMMU: S1, S12, S13, S14, S15, S16, S17, S18, B2. From these DMMU, actual sampling was from areas with very little dredged material. Of this group, DMMU S1, S15 and B2 failed PSDDA guidelines for open-water disposal and the rest were considered suitable.

Though this sampling event may have accurately characterized the proposed dredging prism, there is no way whether to know whether this is the case. Therefore, with insufficient information for decision-making, DMMUs S12, S13, S14, S16, S17 and S18 cannot be considered suitable for open-water disposal.

Dredging Year 2001

<u>Weyerhaeuser Company/Hylebos Wood Debris Group</u>. The Weyerhaeuser Company dredging project was located at the head of Hylebos Waterway, Commencement Bay in a MTCA cleanup area. The following biological testing summary covers the use of best professional judgement exercised by the DMMP agencies during sediment testing for this project.

Standard bioassay testing was conducted on all nine DMMUs within the 56 day biological holding time. **Appendix B** summarizes the solid phase bioassay Quality Control (QC) performance guidelines and also summarizes the solid phase bioassay interpretative guidelines for nondispersive sites, which were used to evaluate the bioassay data discussed below. Reference sediment was collected from Carr Inlet to conduct DMMU specific test sediment comparisons for the three bioassay tests used during the three testing rounds. Amphipod bioassays conducted during Round 1 testing with *Ampelisca abdita* failed to meet the negative control performance standard, but the single DMMU tested with Rhepoxynius abronius met both the negative control and reference performance standards. Amphipod testing during round 2 was problematic, based on performance standard failures for both reference and negative control sediments for both amphipod species (Rhepoxynius abronius and Ampelisca abdita), which resulted in a requirement to retest a subset of the DMMUs tested. Round 3 amphipod resampling/retesting of 8 of 9 Round 1 DMMUs was conducted with *Ampelisca abdita*, and all Round 3 batches met both the negative control and reference sediment performance standards. Round 2 testing of Weyerhauser sediments was accomplished with the echinoderm (*Dendraster excentricus*), and the sediment larval bioassay meet both the negative control performance and reference performance standards. In general, the *Neanthes* growth bioassay met the DMMP performance standards for the negative control and reference sediments during Rounds 1 and 2 and the results will not be discussed further. Summary bioassay results for each DMMU are depicted in **Appendix C** relative to the DMMP nondispersive interpretative guidelines. Bioassay results are discussed below for the amphipod and sediment larval bioassays only.

a) Amphipod Bioassay (Ampelisca abdita, Rhepoxynius abronius). The cumulative results for amphipod bioassay testing conducted over three testing rounds are summarized in Table A-7. The testing resulted in 5 DMMUs among the three testing rounds exhibiting no-hit responses, 1 DMMU with 2-hit responses, and 3 DMMUs with a 1-hit response. DMMU B-3 had 1-hit responses from both the Rhepoxynius abronius and Ampelisca abdita bioassays during Round 1 and Round 1 retest.

- b) Bivalve Larval Bioassay (Dendraster excentricus, Mytilus galloprovincialis). Of the Eight of nine DMMU's⁴ tested, three passed the DMMP nondispersive guidelines for unconfined open-water disposal, and five exhibited a 2-hit responses (Table A-7). One of the three suitable DMMUs exceeded the 2-hit response guidelines, but when statistically compared to the appropriate reference sediment responses, was found to be "not statistically different" (p<0.1), and thus is not scored as a "hit" for regulatory decision-making (e.g., "suitable").
- c) <u>DMMP Bioassay Determination</u>. Overall interpretation of the nine DMMU's tested by the DMMP bioassays demonstrated that five DMMUs exhibited bioassay responses that were suitable for unconfined openwater disposal (UOWD) and four exhibited responses that were unsuitable for UCOWD (Appendix C, Table A-7).
- d) <u>Bioaccumulation Trigger Exceedances</u>. Of the six DMMUs that had BT exceedances for TBT, three passed the DMMP bioassays interpretation guidelines for open-water-unconfined disposal during bioassay testing Rounds 1-3. Therefore, three DMMUs (B-5, B-6, B-7) were subject to bioaccumulation testing during Round 4.

Table A-7. Bioassay interpretation summary³

Amphipod bioassay: (Rhepoxynius abronius & Ampelisca abdita)	Pass	Two-Hit	One-Hit	Total:
Weyerhaeuser Dock DMMUs	5	1	3	9
Sediment Larval Bioassay: (Dendraster excentricus)	Pass	Two-Hit	One-Hit	Total:
Weyerhaeuser Dock DMMUs	3	5	0	8
Neanthes Growth Bioassay: (Neanthes arenaceodentata)	Pass	Two-Hit	One-Hit	Total:
Weyerhaeuser Dock DMMUs	9	0	0	9
DMMP Bioassay Determination:		f Suitable MUs		Unsuitable MUs
Weyerhaeuser Dock DMMUs	Í	5	2	4

The sediment analytical chemical results for the 3 DMMUs that underwent bioaccumulation testing for TBT are presented in **Table A-8**. The results of these sediment analyses indicated that there was often a large

A-23

⁴ The larval bioassay test was not rerun on DMMU B-3 (see **Appendix C**) in Round 2 because this DMMU exhibited Round-1 single-hit responses for both *Rhepoxynius abronius* and *Ampelisca abdita*.

disparity between the initial and resampled/retested analytical results for porewater-TBT. When the initial result exceeded the retested result, the ratio of the two was used to adjust the bioaccumulation tissue concentrations to reflect a "worst case" analytical result.

Table A-8. Ratio of Initial Sediment Porewater-TBT Concentrations to Retested Sediments for Bioaccumulation Testing

DMMU ID	Initial (ug/L)	Retest (ug/L)	Initial/Retest Ratio
B-5	0.67	0.23	<mark>2.91</mark>
B-6	0.33	0.16	<mark>2.06</mark>
B-7	0.17	0.073	<mark>2.33</mark>

Bioaccumulation testing was performed with *Macoma nasuta*, a facultative deposit feeding/suspension feeding bivalve and *Nephtys caecoides*, a burrowing facultative deposit feeding/carnivorous polychaete. The two species were tested together in the same 8-gallon aquaria. The protocol used followed the recent DMMP protocol clarification (Kendall, 2000), which extended the standard bioaccumulation test duration from 28 days to 45 days. Five replicate 8-gallon aquaria were run for the negative control/reference sediments, and for each of the three tested DMMUs.

Tissue concentrations of chemicals-of-concern from the 45-day exposures were compared statistically to the appropriate reference sediment, based on grain size similarity comparisons. As noted above, the calculated ratios of Phase 1 (initial)/Phase II (retest) sediment chemistry were used to adjust the observed tissue concentrations (**Table A-8**). Statistical comparisons of test DMMUs and reference tissue concentrations for the final interpretation "worst case" analyses were based on the adjusted tissue concentrations. The summary tissue chemistry interpretation for each of the measured chemicals is provided in **Table A-9** for each of the 3 DMMUs tested.

The DMMP agencies agreed that comparing statistical differences from reference is a necessary, but not sufficient condition to determine a DMMU unsuitable for open-water disposal. For those DMMUs that were statistically greater than reference, a more in depth evaluation was required to determine the significance of the bioaccumulation that had occurred. This evaluation focused on a) Food and Drug Administration (FDA) Action Levels for Poisonous and Deleterious Substances in Fish and Shellfish for Human Food; b) PSDDA target tissue concentration values for chemicals of concern to human health, and c) ecological residue-effects data from the literature.

There are no FDA guidelines for Tributyltin (TBT).

A target tissue trigger level (TTL), of 3 ppm dry weight of TBT in tissue (0.6

ppm wet weight) was used to evaluate the Weyerhaeuser Dock tissue concentrations. This tissue concentration is protective for growth and reproduction endpoints in polychaetes, crustaceans, bivalves, and most gastropods. However, it might not protect the most sensitive species of meso-and neogastropods against imposex-related sterility. Considering that meso-and neogastropods are rare in Elliott Bay (Appendix D in EVS, 1999) and are thus presumed to be rare in Commencement Bay as well, the DMMP agencies have decided to extrapolate the use of the **TBT trigger level (3 ppm dry weight)** derived for the West Waterway on an interim basis to interpret bioaccumulation data relative to disposal at the Commencement Bay disposal site.

To summarize, the DMMP agencies used the following TTLs to interpret the bioaccumulation test data for the proposed Weyerhaeuser Dock dredging area:

TBT: 3.0 ppm dry weight (dw) as TBT

The agencies used best professional judgement in developing the interpretation guidelines to meet PSDDA disposal site management objectives; achievement of other sediment management objectives will require additional evaluation. The use of this guideline is subject to change for future PSDDA/DMMP projects as additional bioaccumulation data become available.

Each of the three DMMU's was compared to these interpretation guidelines using a one-tailed one-sample t-test (see **Table A-9**). An alpha level (the probability of making a Type I error, rejecting the null hypothesis of no difference between test and reference responses when, in fact, they are not different) of 0.1 was selected for these statistical comparisons by the DMMP agencies to reflect the higher within sample variability, and to increase the power of the test to discriminate between reference and test responses. All three DMMU's were found to be statistically less than the TBT target tissue level (TTL) and thus passed the bioaccumulation test.

 Table A-9. Bioaccumulation Testing Summary for Weyerhaeuser DMMUs tested under DMMP.

							DMMU B-5					
					Macoma nasuta					Nephtys caecoides		
CHEMICAL NAME	Units	Guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference	Statistically different from reference	Statstically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference	Statistically different from reference	Statstically below guideline
TBT ion (as TBT)	ug/kg-dw	3,000	125	363	4.4	yes	yes	28.1	81.9	0.9	yes	yes
							DMMU B-6					
					Macoma nasuta					Nephtys caecoides		
CHEMICAL NAME	Units	Guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference	Statistically different from reference	Statstically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference	Statistically different from reference	Statstically below guideline
TBT ion (as TBT)	ug/kg-dw	3,000	176	364	4.4	yes	yes	37.6	78	0.9	yes	yes
							DMMU B-7					
					Macoma nasuta					Nephtys caecoides		
CHEMICAL NAME	Units	Guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference	Statistically different from reference	Statstically below guideline	DMMU tissue (Initial)	DMMU tissue (adjusted)	Reference	Statistically different from reference	Statstically below guideline
TBT ion (as TBT)	ug/kg-dw	3,000	107	250	4.4	yes	yes	25.2	59	0.9	yes	yes

Port of Anacortes, Cap Sante Marina. Nine of twelve DMMUs tested for the Cap Sante Marina project had screening level exceedances of tributyltin. Additional sediment was collected for bioaccumulation testing based on an addendum to the original sampling plan. Two test composites were created, one representing those samples with TBT detections above 0.30 uG/L and those with detections below 0.30 uG/L but above the SL. TBT exceedances are as follows:

	C1	C2	C6	C7	C8	C9	C10	C11	C12	Comp1	Comp2
TBT in ug/L	0.47	0.20	0.18	0.34	0.29	0.24	0.32	0.27	0.20	0.30	0.29

Bioaccumulation testing was performed with bivalve *Macoma nasuta* and the polychaete *Nephtys caecoides*. The two species were tested together in the same 18-liter glass aquarium. At the time of project initiation, the standard DMMP bioaccumulation protocol called for 28-day test duration. The project proponents agreed to extend the test to 45 days, based on the recommendation of the DMMP agencies. The extended test provides a better approximation of steady-state tissue concentrations for TBT.

Six replicate aquaria (five test replicates and one replicate for steady state monitoring) were run for the two test sediments, the two reference sediments and the negative control. Tissue concentrations from the 45-day exposure were compared to the reference sediments. Initial sediment chemistry was used to adjust the observed tissue concentrations.

The DMMP agencies agreed to use the target tissue level developed for the East Waterway project, 3 ppm dry weight of TBT in tissue, as the value appropriate for the Cap Sante project. Given the limited residue-effects data for the Anacortes area, it was determined that the number calculated for Elliott Bay would be the closest approximation available for making a determination of suitability. The method of calculation and the data supporting this value is documented in the suitability determination for the East Waterway project suitability determination (1999), paragraph 18, and in the "Review of Tissue Residue Effects Data for Tributyltin, Mercury and Polychlorinated Biphenyls", prepared by EVS solutions for the Port of Seattle.

TBT concentrations in tissues from *Macoma* and *Nephtys* exposed to test sediments were significantly less than the target tissue level of 3 ppm dry weight TBT in tissue. TBT tissue concentrations were also compared to reference and no significant differences were observed for both DMMU.

<u>USACE Squalicum Waterway, Bellingham</u>. The Squalicum Waterway is a federal navigation channel in Bellingham, Washington. It is of mixed rank, with

the outer portion ranked moderate and the head of the waterway, near historical industrial use, ranked high. Out of 17 DMMU, 13 had no detected or non-detected chemical exceedances of screening levels (**Table A-10**). Three of those DMMU with SL exceedances (C5, C6 and C7) exhibited borderline exceedances of nickel but no other chemicals of concern. The agencies used best professional judgment for waiving bioassays on these three DMMU, as described below. The fourth DMMU that exceeded screening levels was S2, with high levels of lead found. The agencies defined further testing for S2 before it could be found suitable for open-water disposal, also as described below.

Table A-10. Selected Squalicum Waterway chemistry results.¹

PARAMETER			hemica uidelin BT		C5 (M)	C6 (M)	C7 (M)	S2 (H)
Volume (cubic yards)					15,164	11,564	15,659	1,688
METALS (ppm)	Lead	450		1,200				2,100
	Nickel	140	370	370	140	141	140	
2,4-Dimethylphenol (ppb)		29		210				62

'Table includes all chemicals of concern (COCs) that exceeded PSDDA SL triggers.

No additional COCs exceeded SLs.

1. Nickel Exceedances. Three DMMU (C5, C6 and C7) in the turning basin area of the navigation channel had nickel detected at or close to (140, 141 and 140 ppm, respectively) the SL (140 ppm). Analysis of the data showed that Ni concentrations were localized in the surface portion of the turning basin material, and concentrations decreased with increasing distance from the turning basin. Grain size showed a high negative correlation with Ni concentration (Figure 1). Similar concentrations have been found in the waterway in past sampling. Though SL exceedances usually trigger Tier 3 bioassays before being found suitable for open water disposal, the agencies' suspended this requirement for these DMMU. This "best professional judgement" decision was based on the following lines of evidence:

similar and greater levels of Ni found in the same area before have passed bioassays

there is circumstantial evidence that background levels of Ni in the Bellingham Bay area are high

there are no apparent anthropogenic sources

these DMMU showed no SL exceedances of any other COC




Figure A-1. Ni concentrations vs. percent fines.

2. Lead and 2,4-dimethylphenol Exceedances in S2. Subsurface DMMU S2, in the right berth area, had both an SL exceedance of 2,4-dimethylphenol and an ML exceedance of lead (Table A-10). This level of lead caused much more concern among the agency representatives than did the detections of nickel. To establish an approach for further testing of this DMMU, the agencies looked at the site history, laboratory QA/QC data, and potential human health and environmental effects of lead.

A review of the site history showed that both a plywood company and a boat building business have been located in the area north of the right berthing area since the 1920s. Boat building, of both metal and wooden boats, ceased in the 1960's, though a plywood company continues to operate today. This area was filled to its present shoreline in the 1940s and 50s. At present, the source or extent of subsurface contamination is undetermined.

The analytical laboratory reviewed its procedures and found no indication that the lead measurement was erroneous. There are also potential human as well as environmental effects of lead contamination.

With the above information, the agencies agreed on a plan for further evaluation of S2. The following Tier 3 tests were required before S2 could be considered for open water disposal:

- ?? Acute bioassays with the usual suite of bioassay tests. *Mytilus* should be the species chosen for the larval bioassay due to its sensitivity to metals. Should the sediment fail these tests, it would be unsuitable for open-water disposal.
- 28-day bioaccumulation tests on two species before the sediment could be considered suitable for open-water disposal. Tissue analysis would be for Pb only.
- 2? Should the area be dredged, regardless of disposal area, Z samples would need to be analyzed to make sure that the exposed surface does not exceed Ecology non-degradation standards. Further dredging may be necessary if contamination is shown to go deeper than the proposed dredging prism.
- ?? Bioassay and bioaccumulation tests could be tiered or conducted concurrently.
- 7? Though it was found suitable for open-water disposal, no portion of S1 (overlying S2) should be removed without further testing of subsurface sediments.

Bellingham Cold Storage chose not to pursue further action at this site as part of the current project.

<u>USACE Grays Harbor Maintenance Dredging.</u> Requirements for dredged material characterization in the Grays Harbor Navigation Channel vary in many ways from requirements in other parts of Washington. The channel is dredged annually of over a 1.5 million cubic yards of homogenous, low-ranked material deposited from a large river system into a relatively shallow estuary. The Grays Harbor/Willapa Bay Dredged Material Evaluation Procedures (GHDMEP) outlines a six-year rotation of sampling and testing, with one-third of the channel material tested every two years. The first six year testing cycle was completed in 1999, so the DY 2001 testing began the second cycle. This paragraph describes adaptations to the typical SAP used for this project, and the use of best professional judgment for the choice of sample used for safetynet bioassay testing.

- 1. <u>Programmatic Sampling Plan</u>. Some "lessons learned" during the first round of testing included:
 - ?? keeping track of areas tested between testing events was difficult and confusing, leading to concerns that some areas or problems could be overlooked
 - ?? material to be dredged is concentrated in the upper reaches of the channel, where the possibility of contamination is greatest, and is not easily divided into thirds for testing

To address these concerns, a Programmatic Sampling and Analysis Plan (PSAP) was prepared to outline the sampling approach for the entire six-year cycle. This PSAP looked at historic dredging volumes in various reaches of the navigation channel and devised a strategy for insuring that the sampling adequately represented those volumes. A SAP addendum was also prepared to address sampling issues specific to the DY 2001 sampling and testing event. If needed, addenda will also be prepared for subsequent sampling events during the six-year testing rotation.

2. <u>Bioassay Sample Selection</u>. Although no detected levels of chemicals of concern exceeded SLs, the detection levels of four COCs exceeded the SLs in one composite (C10). As per guidelines, the analytical laboratory did everything possible to bring down detection limits but was unable to in this sample. According to DMMP guidelines, exceedance of SLs solely by detection limits can trigger biological testing. In Grays Harbor, safety-net bioassay testing on at least two DMMU composites is required by the GHDMEP. Tiered testing due to SL exceedances (detected or non-detected) can be included in the two samples chosen for safety-net bioassays.

For several reasons, C10 was NOT one of the DMMUs chosen for safety-net bioassays. The selection process considers grain size (fine-grained sediments are preferred); sediment chemical results, and the proximity of the collected samples to known or potential contaminant sources. For this characterization, there were three general groups of grain sizes found in the composite samples:

- ?? <u>Fine grained</u>: C1 through C6 all had over 70% fines (representing 58% of all characterized sediments)
- ?? <u>Intermediate grained</u>: C7 and C8, with around 50% fines (representing 19% of all characterized sediments)
- ?? Coarse grained: C9, C10 and C11 ranged from 5% to 27% fines (representing 23% of all characterized sediments)

Most of the sediments in the Cow Point area are very fine-grained, and they represent the bulk of material deposited annually from upstream sources and dredged in a given year. Because the fine-grained DMMUs all showed similar chemical results, C5, with 74% fine-grained material, was randomly chosen to represent this group of sediments. The remainder of the sediments included two intermediate grain-sized and three coarser-grained DMMU. The intermediate- and coarser-grained sediments were all from the area upstream of the Cow Pt. turning basin, in the Aberdeen and South Aberdeen Reaches. The coarser upstream sediments are less frequently dredged and are generally closer than the downstream sediments to known or potential contaminant sources, including the Weyerhaeuser pulp mill in Cosmopolis. It was field observations of the area and sediments sampled in C11 that led to its choice for safety-net biological sampling. In addition to the timber-related activities and paper mill, boat maintenance activities were observed in the area of

DMMU 11. The sediments collected for C11 exhibited a marked oily sheen and odors. Though chemical analysis of the coarse grained sediments showed no detected levels of COCs above the SL, C11 was chosen over C10 for safety-net testing because it was considered a higher risk for potential toxicity based on field observations.

APPENDIX B - DY00/01 GUIDELINE VALUES (CHEMISTRY)

CHEMICAL NAME	Units	SL	вт	ML	(SL+ ML)/2
METALS & ORGANOMETALS					
Antimony	mg/kg	150	150	200	175
Arsenic	mg/kg	57	507.1	700	378.5
Cadmium	mg/kg	5.1		14	9.55
Copper	mg/kg	390		1,300	845
Lead	mg/kg	450		1,200	825
Mercury	mg/kg	0.41	1.5	2.3	1.355
Nickel	mg/kg	140	370	370	255
Silver	mg/kg	6.1	6.1	8.4	7.25
Zinc	mg/kg	410	-	3,800	2105
TBT ion (porewater)	ug/L	0.15	0.15		
LPAH					
Naphthalene	ug/kg	2,100		2,400	2250
Acenaphthene	ug/kg	500		2,000	1250
Acenaphthylene	ug/kg	560		1,300	930
Fluorene	ug/kg	540		3,600	2070
Phenanthrene	ug/kg	1,500		21,000	11250
Anthracene	ug/kg	960		13,000	6980
2-Methylnaphthalene	ug/kg	670		1,900	1285
Total LPAHs	ug/kg	5,200		29,000	17100
НРАН					
Fluoranthene	ug/kg	1,700	4600	30,000	15850
Pyrene	ug/kg	2,600		16,000	9300
Benzo(a)anthracene	ug/kg	1,300		5,100	3200
Benzofluoranthenes (b+k)	ug/kg	3,200		9,900	6550
Chrysene	ug/kg	1,400		21,000	11200
Benzo(a)pyrene	ug/kg	1,600	3600	3,600	2600
Indeno(1,2,3-c,d)pyrene	ug/kg	600		4400	2500
Dibenzo(a,h)anthracene	ug/kg	230		1900	1065
Benzo(g,h,l)perylene	ug/kg	670		3200	1935
Total HPAHs	ug/kg	12,000		69,000	40500
CHLORINATED HYDROCARBONS					
1,2,4-Trichlorobenzene	ug/kg	31		64	47.5
1,2-Dichlorobenzene	ug/kg	35	37	110	72.5
1,3-Dichlorobenzene	ug/kg	170	1,241		
1,4-Dichlorobenzene	ug/kg	110	120	120	115
Hexaclorobenzene (HCB)	ug/kg	22	168	230	126
PHTHALATES					0
Bis(2-ethylheyesyl)phthalate	ug/kg	8,300	13,870		4150
Butylbenzylphthalate	ug/kg	970			
Di-n-butylphthalate	ug/kg	5,100	10,200		
Di-n-octylphthalate	ug/kg	6,200			
Diethylphthalate	ug/kg	1,200			
Dimethylphthalate	ug/kg	1,400	1,400		

APPENDIX B - DY00/01 GUIDELINE VALUES (CHEMISTRY)

CHEMICAL NAME	Units	SL	вт	ML	(SL+ ML)/2
PHENOLS					
2-Methylphenol	ug/kg	63		77	70
4-Methylphenol	ug/kg	670		3,600	2135
2,4-Dimethylphenol	ug/kg	29		210	119.5
Pentachlorophenol	ug/kg	400	504	690	545
Phenol	ug/kg	420	876	1,200	810
MISCELANEOUS EXTRACTABLES					
Benzyl alcohol	ug/kg	57		870	463.5
Benzoic acid	ug/kg	650		760	705
Dibenzofuran	ug/kg	540		1,700	1120
Hexachlorobutadiene	ug/kg	29	212	270	149.5
Hexachloroethane	ug/kg	1,400	10,220	14,000	7700
N-Nitrosodiphenylamine	ug/kg	28	130	130	79
VOLATILE ORGANICS					
Ethylbenzene	ug/kg	10	27	50	30
Tetrachloroethene	ug/kg	57	102	210	133.5
Total Zylene (sum of o,m,p)	ug/kg	40		160	100
Trichloroethane	ug/kg	160	1,168	1,600	880
PESTICIDES AND PCBs					
Total DDT	ug/kg	6.9	50	69	37.95
Aldrin	ug/kg	10	37		
alpha-Chlordane	ug/kg	10	37		
Dieldrin	ug/kg	10	37		
Heptachlor	ug/kg	10	37		
gamma-BHC (Lindane)	ug/kg	10			
Total PCBs	ug/kg	130	38 (1)	3,100	1615

⁽¹⁾ mg/kg - carbon normalized

Appendix B. DY00/01 DMMP BIOASSAY PERFORMANCE STANDARDS AND EVALUATION GUIDELINES

Bioassay	Negative Control Performance Standard	Reference Sediment Performance Standard	Dispersive Disposal Site Interpretation Guidelines		Nondispersive Disposal Site Interpretation Guidelines		
			1-hit rule	2-hit rule	1-hit rule	2-hit rule	
Amphipod	$M_C \leq 10\%$	M_R - $M_C \le 20\%$	$\begin{array}{c} M_T \text{ - } M_C > 20\% \\ \text{and} \\ M_T \text{ vs } M_R \text{ SD (p=.05)} \\ \text{and} \end{array}$		$\begin{array}{c} M_T \text{ - } M_C > 20\% \\ \text{and} \\ M_T \text{ vs } M_R \text{ SD (p=.05)} \\ \text{and} \end{array}$		
			M_{T} - M_{R} > 10%	NOCN	$M_T - M_R > 30\%$	NOCN	
Larval	$N_C \div I \ge 0.70$	$N_R \geq N_C \geq 0.65$	$\begin{array}{c} N_{T} \div N_{C} < 0.80 \\ \text{and} \\ N_{T}/N_{C} \ vs \ N_{R}/N_{C} \ SD \ (p=.10) \\ \text{and} \end{array}$		$N_{T} \div N_{C} < 0.80$ and $N_{T}/N_{C} \ vs \ N_{R}/N_{C} \ SD \ (p=.10)$ and		
			$N_R/N_C - N_T/N_C > 0.15$	NOCN	$N_R/N_C - N_T/N_C > 0.30$	NOCN	
Neanthes growth	$\begin{aligned} M_C &\leq 10\% \\ &\text{and} \\ MIG_C &> 0.38 \end{aligned}$	$\begin{aligned} M_R &\leq 20\% \\ & \text{and} \\ MIG_R \div MIG_C &\geq 0.80 \end{aligned}$	$\begin{aligned} MIG_T \div MIG_C &< 0.80 \\ &\text{and} \\ MIG_T \text{ vs } MIG_R \text{ SD } (p\text{=}.05) \\ &\text{and} \end{aligned}$		$\begin{aligned} \text{MIG}_{\text{T}} & \div \text{MIG}_{\text{C}} < 0.80 \\ & \text{and} \\ & \text{MIG}_{\text{T}} \text{ vs MIG}_{\text{R}} \text{ SD (p=.05)} \\ & \text{and} \end{aligned}$		
			$MIG_{T}/MIG_{R} < 0.70$	NOCN	$MIG_T/MIG_R < 0.50$	$MIG_{T}/MIG_{R} < 0.70$	

 $M = mortality, N = normal \ survivors, I = initial \ count, MIG = mean \ individual \ growth \ rate \ (mg/individual/day) \\ SD = statistically \ different, NOCN = no \ other \ conditions \ necessary, N/A = not \ applicable$

Subscripts: R = reference sediment, C = negative control, T = test sediment

APPENDIX C – LEGEND

С		
S	=	reported concentration exceeds screening level
SB	=	reported concentration exceeds screening level and bioaccumulation trigger
M	=	reported concentration exceeds maximum level
BM	=	reported concentration exceeds bioaccumulation trigger and maximum level
(U)	=	detection limit exceeds either screening level, bioaccumulation trigger, or
		maximum level
(B)	=	analyte detected in corresponding blank
(E)	=	estimate
(J)	=	detected between the SDL and the CRDL
(UJ)	=	analyte not detected above the sample quanititation limit; however the
		reported quantitation limit is approximate
(D)	=	compound required a dilution as a result of the matrix or the sample
		concentration
(M)	=	estimated value of analyte found and confirmed by analyst, but with low
		spectral match
(N)	=	estimate based on presumptive evidence
(G)	=	estimate is greater than value shown
(Y)	=	raised non-detect due to matrix interferences
NA	=	not analyzed
2H	=	a hit under two-hit interpretation guideline
1H	=	a hit under one-hit interpretation guideline
PASS	=	test sediment passes DMMP guidelines for open-water unconfined disposal
FAIL	=	test sediment fails DMMP guidelines for open-water unconfined disposal
FAIL(C)	=	DMMU found unsuitable for open-water disposal in the absence of
		bioaccumulation and/or Tier IV testing data
(BPJ)	=	best professional judgement applied to suitability determination
L	=	the highest reported concentration was below SL
LM	=	the highest reported concentration was between SL and $(SL + ML)/2$
M	=	the highest reported concentration was between (SL + ML)/2 and ML
Н	=	the highest reported concentration exceeded ML
H*	=	the sediment rank is based on biological testing results

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT: Eastwaterway Project (Phase 2)	S14 H	S15 H	S16
METALS & ORGANOMETALS Antimony Antimony Antimony Arsenic Cadmium	<u> </u>		Н
Arsenic Cadmium Signature			
Cadmium			
Copper			
Lead			
Mercury S S S S S S S S	S	S	S
Nickel			
Silver			
Zinc			
TBT (on (porewater) SB			
LPAH Naphthalene	1		
Naphilladerie Acenaphthene	1	1	
Acenaphthylene			
Fluorene			
Phenanthrene			
Anthracene			
2-Methylnaphthalene	1	1	
Total LPAHs HPAH			
Fluoranthene	1		
Pyrene S S	1	1	
Benzo(a)anthracene			
Benzofluoranthenes (b+k)			
Chrysene	1		
Benzo(a)pyrene	 	<u> </u>	
Indeno(1,2,3-c,d)pyrene	 	 	
Dibenzo(a,h)anthracene Benzo(g,h,l)perylene	1		
Delizorgaria/per yiene			
CHLORINATED HYDROCARBONS			
1,2,4-Trichlorobenzene			
1,2-Dichlorobenzene			
1,3-Dichlorobenzene			
1,4-Dichlorobenzene	1	1	
Hexaclorobenzene (HCB) PHTHALATES			
Bis(2-ethylhyesyl)phthalate			
Buylbenzylphthalate Buylbenzylphthalate			
Di-n-buty/phthalate			
Di-n-octylphthalate			
Diethylphthalate Diethylphthalate			
Dimethylphthalate			
PHENOLS 2-Methylphenol			
Z-metryphenol 4-Metryphenol			
2.4-Dimethylphenol 2.4-Dimethylphenol			
Pentachlorophenol			
Phenol			
MISCELANEOUS EXTRACTABLES			
Benzyl alcohol Benzyl alcohol			
Benzoic acid	+	 	
Dibenzofuran Hexachlorobutadiene	1	 	
riexaciniototicalerie Hexachloroptiane	1		
N-Nitrosodiphenylamine	1		
VOLATILE ORGANICS UCLATIVE ORGANICS			
Ethylbenzene			
Tetrachloroethene	1	ļ	
Total Zylene (sum of o,m,p)	1	 	
Trichloroethane PESTICIDES AND PCBs	+	 	
Total DDT	S (UJ)	S (UJ)	SB (UJ)
Aldrin S(0) S(0) S(0) S(0) S(0)	S (U)	C (00)	S (UJ)
alpha-Chlordane			
Dieldrin S (U) S (U) S (U) S (U)	S (U)		S (U)
Heptachlor S (U)	S (U)		S (U)
gamma-BHC (Lindane)	65	<u> </u>	00
Total PCBs S SB	SB	S	SB
BIOASSAYS Amphined	1		
Amphipod Sediment Larval (Bivalve/Echinoderm) 1H 2H	1	2H	
Secument Latva (produce Culmidatini) 11 21 21 21 21 21 21 21 21 21 21 21 21	1	211	
Bioassay: (Pass/Faii) FAIL PASS PASS PASS PASS PASS PASS PASS PAS	PASS	PASS	PASS
BTs eyesceeded: yes no no yes	yes	no	yes
Bioaccumulation test conducted: no yes yes yes yes yes yes yes yes yes no yes	yes		yes
Bioaccumulation (Pass/Fail): PASS PASS PASS PASS PASS PASS PASS FAIL PASS	PASS		FAIL
ML Rule exceeded: no	no	no	no
OVERALL PASS/FAIL: PASS PASS PASS PASS PASS PASS PASS PAS	PASS	PASS	FAIL
HIGHEST RANKING: H* L L LM LM LM LM M LM H* H* LM	M	LM	H*

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PARTICIDAD 19	PROJECT:							Fastwate	rway Pr	niect (Pha	se 2)						
## CAME AS COMMONDED AS A STATE OF THE PROPERTY OF THE PROPERT	DMMU ID:							S23	S24	S25	S26						
March Marc	METALS & ORGANOMETALS	-17	п			- 11	- 11		- 17		17	17	п		п		
Carrier	Antimony																
Company									c	c							
Second S									3	3							
SSES SINGLE STATE OF THE STATE	Lead																
Sheer The Comment of	Mercury	S						S	S						S		
Trial processors																	
TRY IN CONTROLLED 10 10 10 10 10 10 10 1									S	S							
Page	TBT ion (porewater)	SB (J)				SB (M)		SB (J)			SB (MB)					SB (B)	
Accordation	LPAH																
The propriets The propriet									Q	9							
Riscord	Acenaphthylene								J								
Andersones	Fluorene																
Control Libration									S	S							1
Trigon Parks										S							
PANE	Total LPAHs																
Figure Service	HPAH																
	Fluoranthene									S							
Secreption primary from the Company of the Company																	
Compares	Benzofluoranthenes (b+k)																
Description	Chrysene																
Demorphis plantenageme	Benzo(a)pyrene																
Bearough Jupeysive																	
Troch HPANS																	
	Total HPAHs																
AD-Chilorobervaries																	
Heacetonochemore (HCS)	1,3-Dichlorobenzene																
######################################	1,4-Dichlorobenzene																
Bid2 - ethylogenylphthalate																	
Sulphensylphenialate																	
District physical	Butylbenzylphthalate																
Dietrylephalate	Di-n-butylphthalate																
Dimethyphenical																	
### PHENOLS ### Afterhylphenol #																	
### Afterlightenord ### Af	PHENOLS																
2.4-Dimetrylyphenol	2-Methylphenol																
Pentachiorophenol																	
Phenol																	
Benzyal alcohol	Phenol																
Benzoic acid																	
Dibenzofuran																	
Hexachloroutadiene																	
N-Nitrosodiphenylamine	Hexachlorobutadiene																
	Hexachloroethane																
Ethylbenzene																	
Total Zylene (sum of o,m,p)	Ethylbenzene																
Control Cont	Tetrachloroethene																
Destriction	Total Zylene (sum of o,m,p)																
Total DDT																	
Aldrin S (U)	Total DDT	SB (UJ)		S (U)	S (U)	S (UJ)		BM (U)	BM (J)	BM (U)	S (UJ)	S (UJ)	S (J)	S (J)	S (J)		
Dieldrin S (U)	Aldrin				, , ,			S (U)									
S (U) S (U	alpha-Chlordane	C (LI)				C (LI)		0/10			C /I IV						
Sediment						S (U)		S (U)			S (U)						
SB	gamma-BHC (Lindane)	J (J)							- (O)								
Amphipod 2H 2H 2H 2H 2H 2H 1H 2H	Total PCBs	SB		SB	S	SB		BM	SB	S	S	S	S		S		
Sediment Larval (Bivalve/Echinoderm)	BIOASSAYS Amphiped	211															
Nearthes Growth 2H				2H	2H	2H	2H	2H	1H	2H	2H		2H		2H	2H	2H
Bioassay: (Pass/Fail)	Neanthes Growth																
Bioaccumulation test conducted: no yes yes yes no no no no yes yes	Bioassay: (Pass/Fail)		PASS	PASS		PASS	PASS	PASS				PASS	PASS	PASS	PASS	PASS	PASS
Bioaccumulation (Pass/Fail):	BTs eyesceeded:		no		no		no					no	no	no	no		no
ML Rule exceeded:		no							no	no	no				}		
OVERALL PASS/FAIL: FAIL PASS PASS FAIL PASS PASS FAIL PASS PASS FAIL PASS PASS PASS FAIL PASS	ML Rule exceeded:	no	no		no		no		yes	no	no	no	no	no	no		no
HIGHEST RANKING: H' L LM H' LM L H' H' H' LM LM LM LM H' L	OVERALL PASS/FAIL:	FAIL	PASS	PASS	FAIL	PASS	PASS	FAIL	FAIL (c+b)	FAIL	FAIL	PASS	PASS	PASS	PASS	FAIL	PASS
	HIGHEST RANKING:	H*	L	LM	H*	LM	L	H*	H*	H*	H*	LM	LM	LM	LM	H*	L

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT: DMMU ID: Testing Rank:	S33 H	S34 H	S35 H	S36 H	S37 H	S38 H	Eastwate S39 H	erway Pro S40 H	oject (Pha S41 H	ase 2) S42 H	S43 H	S44 H	S45 H	S46 H	S47 H	S48 H
METALS & ORGANOMETALS																
Antimony																
Arsenic		<u> </u>	<u> </u>				 	 					<u> </u>	<u> </u>		
Cadmium	-	ļ			S	\vdash	$\vdash \vdash$	 					 	 		
Copper Lead					 								+	+		
Mercury	S		S	S	S							S	S			S
Nickel																
Silver													1	1		BM
Zinc					S				S							S
TBT ion (porewater)			SB (MB)	SB (MB)		SB (M)	SB (M)	SB (M)	SB (M)		SB (MB)			SB	SB	SB (M)
LPAH																
Naphthalene		ļ			S		$\vdash \vdash \vdash$			<u> </u>		<u> </u>	 	 		
Acenaphthene		<u> </u>	S		S			 					 	 		
Acenaphthylene Fluorene			S		S			—								
Phenanthrene			S		S											
Anthracene		† ·											1	1		†
2-Methylnaphthalene					S								1	1		S
Total LPAHs					S											
НРАН																
Fluoranthene			S	S	S	igsquare	igsquare	└	igsqcup	——	igsquare	<u> </u>	└	<u> </u>		S
Pyrene	 	<u> </u>	S			igsquare			ldot		\longmapsto	—	<u> </u>	<u> </u>		S
Benzo(a)anthracene	 	 	-	 		$\vdash \vdash \vdash$	$\vdash \vdash \vdash$	 		 		├──	├	 		1
Benzofluoranthenes (b+k) Chrysene	 	┼──		$\vdash \vdash \vdash$		$\vdash \vdash$	$\vdash \vdash \vdash$		\vdash		\vdash	\vdash	₩	₩	-	S(J)
Benzo(a)pyrene	 	\vdash	-	\vdash		\vdash	\vdash	\vdash	 	\vdash	\vdash	\vdash	\vdash	\vdash		3(J)
Indeno(1,2,3-c,d)pyrene	l	 	\vdash									-	 	 	1	S (J)
Dibenzo(a,h)anthracene	i							[S (J)
Benzo(g,h,l)perylene																S (J)
Total HPAHs			S													S (J)
CHLORINATED HYDROCARBONS																
1,2,4-Trichlorobenzene																
1,2-Dichlorobenzene		ļ!			lacksquare	<u> </u>							<u> </u>	<u> </u>		
1,3-Dichlorobenzene			ļ		—	igwdown	 	 	-				-	-		1
1,4-Dichlorobenzene		 		S	-	\vdash	\vdash	— —	-				 	 		
Hexaclorobenzene (HCB) PHTHALATES					—		\vdash	-					 	 		++
Bis(2-ethylheyesyl)phthalate					T								1	1		S (J)
Butylbenzylphthalate			 													<u> </u>
Di-n-butylphthalate													<u> </u>			
Di-n-octylphthalate																
Diethylphthalate																
Dimethylphthalate		<u> </u>											<u> </u>	<u> </u>		
PHENOLS						igsquare										
2-Methylphenol	0		ļ			igwdown	 	 	-				 	-		1
4-Methylphenol	S S				-	\vdash	\vdash	\vdash	 		—		-	-		++
2,4-Dimethylphenol Pentachlorophenol	5	1				\vdash		—								++
Phenol																
MISCELANEOUS EXTRACTABLES		 											<u> </u>	<u> </u>		
Benzyl alcohol																
Benzoic acid																
Dibenzofuran																
Hexachlorobutadiene					<u>_</u>	igsquare	igsquare	└	igsqcup	——	igsquare	<u> </u>	└	<u> </u>		┰
Hexachloroethane		 			\vdash	igsquare	igwdapsilon				<u> </u>	 	 	 		\perp
N-Nitrosodiphenylamine	 	 	-	$\vdash \!$		$\vdash \vdash \vdash$	$\vdash \vdash \vdash$		\vdash		\vdash	-	 	 		├
VOLATILE ORGANICS Ethylbenzene	1	 			S	\vdash	$\vdash \vdash$		\vdash	\vdash	\vdash	\vdash	\vdash	\vdash	1	+
Tetrachloroethene	 	\vdash	-	 	3	$\vdash \vdash$	\vdash	$\overline{}$	\vdash	\vdash	\vdash	\vdash	\vdash	\vdash	 	
Total Zylene (sum of o,m,p)	1		 		S (M)							—	 	 	1	
Trichloroethane	l				C (IVI)								<u> </u>	<u> </u>		
PESTICIDES AND PCBs																
Total DDT			SB (UJ)	BM (UJ)	BM (J)										S (UJ)	S (J)
Aldrin			S (U)	S (U)	S (U)	S (U)										
alpha-Chlordane					S (U)	S (U)			$ldsymbol{ldsymbol{eta}}$	$ldsymbol{ldsymbol{eta}}$				$ldsymbol{ldsymbol{ldsymbol{eta}}}$		
Dieldrin	ļ	<u> </u>		SB (U)	S (U)	S (U)	igwdap	<u> </u>					ــــــ	ــــــ		S (U)
Heptachlor	.	 	S (U)	S (U)	S (U)	S (U)	$\vdash \vdash$		\vdash			⊢—	 	 	-	S (U)
gamma-BHC (Lindane)	 		CD.	DM	DM	CD.		 	-	 	—		 			CD.
Total PCBs BIOASSAYS	!	S	SB	ВМ	BM	SB	S	—	S			S	┼	S	S	SB
Amphipod]			ł		i l	i '				ĺ	2H	1		
Sediment Larval (Bivalve/Echinoderm)	2H	1H	2H	1H	1H	1H	2H	2H	2H	2H	2H	2H	2H	2H	2H	1H
Neanthes Growth			2H	1H	1H	1H										2H
Bioassay: (Pass/Fail)	PASS	FAIL	FAIL	FAIL	FAIL	FAIL	PASS	PASS	PASS	PASS	PASS	PASS	FAIL	PASS	PASS	FAIL
BTs eyesceeded:	no	no	yes	yes	yes	yes	yes	yes	yes	no	yes	no	no	yes	yes	yes
Bioaccumulation test conducted:			no	no	no	no	yes	yes	yes		yes			yes	yes	no
Bioaccumulation (Pass/Fail):							PASS	PASS	PASS		PASS			PASS	PASS	
ML Rule exceeded:	no	no	no	yes	yes	no	no	no	no	no	no	no	no	no	no	no
	PASS	FAIL	FAIL	FAIL (c+b)	FAIL (c+b)	FAIL	PASS	PASS	PASS	PASS	PASS	PASS	FAIL	PASS	PASS	FAIL
OVERALL PASS/FAIL: HIGHEST RANKING:	LM	H*	H*	H*	H*	H*	LM	LM	LM	L	LM	LM	H*	LM	LM	H*

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT: DMMU ID: Totiling Registration	S49	S50	S51	S52	S53	S54	Eastwa S55	S56	S57	S58	S59	S60	D1	D2	D3	D4
Testing Rank: METALS & ORGANOMETALS	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н
Antimony																
Arsenic																
Cadmium			S													
Copper																
Lead	0	0	0													-
Mercury	S	S	S													
Nickel Silver																\vdash
Zinc	S	S														
TBT ion (porewater)	SB (MB)	SB (B)		SB (M)					SB (MB)							
LPAH	- C ()	G (()		G = ()					- (····=)							
Naphthalene																
Acenaphthene		S	S													
Acenaphthylene																
Fluorene		S	S													
Phenanthrene		S	S													igwdot
Anthracene		S														$\vdash \vdash$
2-Methylnaphthalene		S	S													\vdash
Total LPAHs HPAH		S	S		1	1	1		 	1	1		1			$\vdash \vdash$
Fluoranthene		SB	S		 	 	 		S	 	 		 			\vdash
Pyrene		S	S		1	1	1		S	1	1		1			\Box
Benzo(a)anthracene		S	S		1		1				1		1			
Benzofluoranthenes (b+k)																
Chrysene		S	S													
Benzo(a)pyrene		S														
Indeno(1,2,3-c,d)pyrene		S														
Dibenzo(a,h)anthracene		S (J)			ļ	ļ	ļ			ļ	ļ		ļ			igwdown
Benzo(g,h,l)perylene		S (J)														\vdash
Total HPAHs CHLORINATED HYDROCARBONS		S (J)	S													\vdash
1,2,4-Trichlorobenzene																\vdash
1,2-Dichlorobenzene																\vdash
1,3-Dichlorobenzene																
1,4-Dichlorobenzene																
Hexaclorobenzene (HCB)																
PHTHALATES																
Bis(2-ethylheyesyl)phthalate																
Butylbenzylphthalate																
Di-n-butylphthalate																—
Di-n-octylphthalate																\vdash
Diethylphthalate																\vdash
Dimethylphthalate PHENOLS																
2-Methylphenol																
4-Methylphenol																
2,4-Dimethylphenol																
Pentachlorophenol																
Phenol																
MISCELANEOUS EXTRACTABLES																
Benzyl alcohol																
Benzoic acid																—
Dibenzofuran			S													—
Hexachloropthone				 	 	 	 		-	 	 		 			\vdash
Hexachloroethane N-Nitrosodiphenylamine				 	1	1	1		 	1	1		1			
VOLATILE ORGANICS					1	1	1			1	1		1			\Box
Ethylbenzene					l	l	l			l	l		l			
Tetrachloroethene																
Total Zylene (sum of o,m,p)																
Trichloroethane																
PESTICIDES AND PCBs					ļ	ļ	ļ			ļ	ļ		ļ			igsquare
Total DDT	S (UJ)	S (UJ)	BM (J)		 		 			 	 		 			\vdash
Aldrin		S (U)	S (U)		 	 	 			 	 		 			$\vdash \vdash$
alpha-Chlordane	C (LI)		S (U)		 	0/11	<u> </u>			 	 		 			$\vdash \vdash$
Dieldrin Heptachlor	S (U)		SB (U) S (U)		 	S (U)			-	 	 		 			\vdash
gamma-BHC (Lindane)			3 (U)		1	1	1		 	1	1		1			
Total PCBs	SB	SB	BM	S	l	S	l		S	S	l	S	l			\vdash
BIOASSAYS	CD	35			1		1				1		1			$\overline{}$
Amphipod				1	1		1			1	1		1			i l
Sediment Larval (Bivalve/Echinoderm)	2H	2H	1H	2H	2H	2H		2H	2H		2H	2H			2H	2H
Neanthes Growth			1H													
Bioassay: (Pass/Fail)	PASS	PASS	FAIL	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS
BTs eyesceeded:	yes	yes	yes	yes	no	no	no	no	yes	no	no	no	no	no	no	no
Bioaccumulation test conducted:	yes	yes	no	yes					yes							igsquare
Bioaccumulation (Pass/Fail):	PASS	PASS		PASS		 	 		PASS		 		 			$\vdash \vdash$
ML Rule exceeded:	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no
OVERALL PASS/FAIL: HIGHEST RANKING:	PASS	PASS	FAIL (c+b)	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS
	LM	LM	H*	LM	L	LM	L	L	LM	LM	L	LM	L	L	L	L

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT:	: D5	D6	D7	D7a	D7b	D7c	D8	D9	D10	D11	D12	D13	D14	D15	D16
Testing Rank: METALS & ORGANOMETALS	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	H	Н
Antimony															
Arsenic															
Cadmium															
Copper				ļ			ļ	ļ				S			
Lead							<u> </u>			 		0	-		
Mercury	S	<u> </u>	S	BM	SB	S	-					S	S		
Nickel Silver		ļ				<u> </u>	 	 				SB			<u> </u>
Zinc	1					S						SB			
TBT ion (porewater)								 							
LPAH								1							
Naphthalene															
Acenaphthene															
Acenaphthylene															
Fluorene															
Phenanthrene		<u> </u>				L	<u> </u>								
Anthracene		<u> </u>		<u> </u>						<u> </u>					
2-Methylnaphthalene	⊢	 	\longmapsto			<u> </u>	——	 	1	$\vdash \vdash$					
Total LPAHs HPAH		┼──	$\vdash \vdash \vdash$		 	<u> </u>			 	$\vdash \vdash$					
Fluoranthene		 	$\vdash \vdash \vdash$		 		 	 	\vdash	$\vdash \vdash$					
Pyrene		 			\vdash			 	-						
Benzo(a)anthracene								 							
Benzofluoranthenes (b+k)	<u> </u>														
Chrysene		†													
Benzo(a)pyrene															, .
Indeno(1,2,3-c,d)pyrene															
Dibenzo(a,h)anthracene															
Benzo(g,h,l)perylene			$ldsymbol{ldsymbol{eta}}$												
Total HPAHs															
CHLORINATED HYDROCARBONS		<u> </u>				L	<u> </u>								
1,2,4-Trichlorobenzene							<u> </u>	<u> </u>		igsquare					
1,2-Dichlorobenzene		ļ					 			\vdash					
1,3-Dichlorobenzene		ļ					 			\vdash					
1,4-Dichlorobenzene		ļ				<u> </u>	 	 							<u> </u>
Hexaclorobenzene (HCB) PHTHALATES		ļ				<u> </u>	 	 							<u> </u>
Bis(2-ethylheyesyl)phthalate	_				\vdash		—								
Butylbenzylphthalate	1														
Di-n-butylphthalate	1														
Di-n-octylphthalate															
Diethylphthalate		† ·						<u> </u>							
Dimethylphthalate															
PHENOLS															
2-Methylphenol															
4-Methylphenol															
2,4-Dimethylphenol				S (U)	S (U)	S (U)									
Pentachlorophenol		<u> </u>				L	<u> </u>								
Phenol		<u> </u>		<u> </u>			<u> </u>			<u> </u>					
MISCELANEOUS EXTRACTABLES		ļ					 			\vdash					
Benzyl alcohol	_		——	S (U)	S (U)	S (U)		ļ		$\vdash \vdash \vdash$					
Benzoic acid Dibenzofuran		 	$\vdash \vdash \vdash$		 			 	\vdash	$\vdash \vdash \vdash$					
Hexachlorobutadiene		\vdash	\vdash		 			 	-	\vdash					
Hexachloroethane								†							
N-Nitrosodiphenylamine		1													
VOLATILE ORGANICS															, .
Ethylbenzene															
Tetrachloroethene															
Total Zylene (sum of o,m,p)			$ldsymbol{ldsymbol{eta}}$												
Trichloroethane					igsquare	ļ	<u> </u>	<u> </u>		igsquare					<u> </u>
PESTICIDES AND PCBs	—	<u> </u>						 							<u> </u>
Total DDT	⊢—	 	BM (UJ)	SB (Y)	SB (Y)	BM (Y)	<u> </u>	 	S (U)	igwdown		S (UJ)	S (J)		
Aldrin		 	00 (11)	<u> </u>	₩	<u> </u>		 		$\vdash \vdash$					
alpha-Chlordane		 	SB (U)		 	 	-	 		├			0 /11/		
Dieldrin Heptachlor		 	SB (U)		 		 		<u> </u>	├			S (U)		
gamma-BHC (Lindane)		\vdash	\vdash		 			 	-	\vdash					
Total PCBs	S		BM	S	SB	SB		S	S	 		S	SB		
BIOASSAYS		 	D.WI		CD	CD						3	JD		\vdash
Amphipod	1	1	2H	1H	1H	1H									
Sediment Larval (Bivalve/Echinoderm)		†		2H	2H	1H			2H		2H	1H	1H	2H	2H
Neanthes Growth	<u> </u>		1H		1H				2H				1H	2H	
Bioassay: (Pass/Fail)	PASS	PASS	1H FAIL	1H FAIL	FAIL	1H FAIL	PASS	PASS	FAIL	PASS	PASS	1H FAIL	FAIL	FAIL	PASS
BTs eyesceeded:	no	no	yes	yes	yes	yes	no	no	no	no	no	yes	yes	no	no
Bioaccumulation test conducted:			no	no	no	no						no	no		
Bioaccumulation (Pass/Fail):															
ML Rule exceeded:	no	no	yes	no	no	no	no	no	no	no	no	no	no	no	no
OVERALL PASS/FAIL: HIGHEST RANKING:	PASS LM	PASS L	FAIL (c+b) H*	FAIL H*	FAIL H*	FAIL H*	PASS L	PASS LM	FAIL H*	PASS L	PASS L	FAIL H*	FAIL H*	FAIL H*	PASS L

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT: DMMU ID: Testing Rank:	D17 H	D18 H	D19 H	D20 H	D21 H	D22 H	D23 H	Eastwate D24 H	erway Pro D25 H	D ject (Ph D26 H	D27	D28 H	D29 H	D30 H	D31 H	D32 H
METALS & ORGANOMETALS																
Antimony																
Arsenic																
Cadmium																
Copper																
Lead Mercury										S						—
Nickel										0						
Silver																
Zinc										1						
TBT ion (porewater)																
LPAH																
Naphthalene																
Acenaphthene																
Acenaphthylene																
Fluorene																
Phenanthrene																
Anthracene 2-Methylnaphthalene																
Total LPAHs																
HPAH																
Fluoranthene																
Pyrene																
Benzo(a)anthracene																
Benzofluoranthenes (b+k)																
Chrysene																
Benzo(a)pyrene																
Indeno(1,2,3-c,d)pyrene																
Dibenzo(a,h)anthracene		ļ					ļ									
Benzo(g,h,l)perylene																
Total HPAHs		-					-									
CHLORINATED HYDROCARBONS 1.2.4-Trichlorobenzene																
1,2,4-1 richiorobenzene 1,2-Dichlorobenzene																—
1,3-Dichlorobenzene																
1,4-Dichlorobenzene																
Hexaclorobenzene (HCB)																
PHTHALATES																
Bis(2-ethylheyesyl)phthalate																
Butylbenzylphthalate																
Di-n-butylphthalate																
Di-n-octylphthalate																
Diethylphthalate																
Dimethylphthalate																<u> </u>
PHENOLS																
2-Methylphenol 4-Methylphenol																—
2,4-Dimethylphenol					S											
Pentachlorophenol					3											
Phenol																
MISCELANEOUS EXTRACTABLES																
Benzyl alcohol																
Benzoic acid																
Dibenzofuran																
Hexachlorobutadiene																oxdot
Hexachloroethane							<u> </u>									
N-Nitrosodiphenylamine		 					 									\vdash
VOLATILE ORGANICS		 					 						-		-	\vdash
Ethylbenzene Tetrachloroethene		1					1									\vdash
Total Zylene (sum of o,m,p)																\vdash
Trichloroethane																
PESTICIDES AND PCBs																
Total DDT																S (U)
Aldrin																` '
alpha-Chlordane																
Dieldrin																
Heptachlor																
gamma-BHC (Lindane)																
Total PCBs	S	S	S													SB
BIOASSAYS		1					1									
Amphipod																
Sediment Larval (Bivalve/Echinoderm)	2H	2H	2H	2H		2H		2H	2H							
Neanthes Growth	DAGG	DAGG	DAGG	DAGG	DAGG	DAGG	2H	DAGG	1H FAIL							
Bioassay: (Pass/Fail)	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS								
BTs eyesceeded:	no	no	no	no	no	no	no	no	yes							
Bioaccumulation test conducted: Bioaccumulation (Pass/Fail):		-	-				-		-							no
ML Rule exceeded:	no	no	no	no	no	no	no	no	no							
OVERALL PASS/FAIL:	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	FAIL							
HIGHEST RANKING:	LM	LM	LM	L L	LM	L	L	L	L	LM	L	L	L	L	L	H*
	LIVI	-141	-141	_	-141		<u> </u>		-	-/41				<u> </u>		

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

Testing Rank: H	H H	H	H	H	S S S	Н	н
Antimony Arsenic Cadmium Copper Lead Mercury Nickel Silver Zinc TBT ion (porewater) LPAH Naphtalene Acenaphthene Acenaphthylene Fluorene Phenanthrene Anthracene 2-Methylnaphthalene Total LPAHS HPAH Fluoranthene Pyrene Benzo(a)anthracene Benzoflouranthenes (b+k) Chrysene	S		S		S		
Cadmium S Copper	S		S		S		
Copper Lead	S	\$	S		S		
Lead Mercury Mickel S Silver S Zinc S TBT ion (porewater) S LPAH Image: Company of the comp	S	\$	S		S		
Mercury S S S S S Nickel Silver Silver S S S S S S S S S	S		S		BM		
Nickel Silver Zinc S TBT ion (porewater) S LPAH Image: Control of the control	S	S	S				
Silver Zinc	S	S	S				
Zinc S S	S	\$	S		S		
LPAH Naphthalene Nacenaphthene	S	S	S				
Naphthalene Acenaphthylene Acenaphthylene ————————————————————————————————————	S	S	S				
Acenaphthene Acenaphthylene Acenaphthylene Image: Common state of the com	S	S	S				
Acenaphthylene Fluorene Phenanthrene ————————————————————————————————————	S	S	S				
Fluorene	S	S	S				
Phenanthrene Anthracene 2-Methylnaphthalene S Total LPAHS Total LPAHS HPAH Fluoranthene Pyrene Benzo(a)anthracene Benzofluoranthenes (b+k) Chrysene	S	S	S				
Anthracene 2-Methylnaphthalene S Total LPAHS HPAH Fluoranthene Pyrene Benzo(a)anthracene Benzo(lovranthenes (b+k) Chrysene	S	S	S				
2-Methylnaphthalene S Total LPAHS HPAH HPAH HIuoranthene Pyrene Benzo(a)anthracene Benzofluoranthenes (b+k) Chrysene Chrysene Inches (b+k)	S	S	S				
Total LPAHS	S	S	S				
HPAH Fluoranthene Pyrene Benzo(a)anthracene Benzofluoranthenes (b+k) Chrysene Chrysene Benzofluoranthenes (b+k) Benzofluoranthenes (b+k) Chrysene Benzofluoranthenes (b+k) Benzofluoranthe	S	S	S				t
Fluoranthene	S	S	S				1 ,
Pyrene Benzo(a)anthracene Benzofluoranthenes (b+k) Chrysene		S	S				
Benzo(a)anthracene Benzofluoranthenes (b+k) Chrysene					S		
Chrysene		l l					
Benzo(a)pyrene							
				-		-	₩
Indeno(1,2,3-c,d)pyrene			S		 	-	+
Dibenzo(a,h)anthracene Benzo(g,h,l)perylene	-		5		 	1	\vdash
Total HPAHs			S				
CHLORINATED HYDROCARBONS							
1,2,4-Trichlorobenzene							
1,2-Dichlorobenzene							
1,3-Dichlorobenzene							
1,4-Dichlorobenzene							
Hexaclorobenzene (HCB)							
PHTHALATES							
Bis(2-ethylheyesyl)phthalate							-
Butylbenzylphthalate Di-n-butylphthalate							1
Di-n-octylphthalate							†
Diethylphthalate							
Dimethylphthalate							1
PHENOLS							
2-Methylphenol S (U)							
4-Methylphenol							
2,4-Dimethylphenol S (U)							
Pentachlorophenol S (U)							
Phenol							
MISCELANEOUS EXTRACTABLES							-
Benzyl alcohol S (U) Benzoic acid S (U)							
Dibenzofuran					 	-	+
Hexachlorobutadiene S (U)							
Hexachloroethane						1	
N-Nitrosodiphenylamine S (U)							
VOLATILE ORGANICS							
Ethylbenzene							
Tetrachloroethene					<u> </u>		↓
Total Zylene (sum of o,m,p)				-		-	₩
Trichloroethane PESTICIDES AND PCBs					-	-	
PESTICIDES AND PCBS	S	S (J)	SB (J)	S	S	S	
Aldrin S (U) S (U) S (U)	3	S (3)	OD (J)	3	3	3	+
alpha-Chlordane S (U) S (U)				1		1	\vdash
Dieldrin S (U) S (U) S (U)							
Heptachlor S (U)							
gamma-BHC (Lindane)							
Total PCBs SB BM BM SB S	S	S	S		S		
BIOASSAYS							
Amphipod 2H							
	2H	1H	1H	1H	1H	2H	
Neanthes Growth 1H 1H 1H 1H 1H 2H	MACC - DAGC	E 0 11	E4.11	E A III	E 6.11	DAGG	DAGG
	PASS PASS	FAIL	FAIL	FAIL	FAIL	PASS	PASS
	no no	no	no	np	yes	no	no
Bioaccumulation test conducted: no no no no no Bioaccumulation (Pass/Fail):	+				no	1	
	no no	no	no	no	no	no	no
	PASS PASS	FAIL	FAIL	FAIL	FAIL	PASS	PASS
	LM LM	H*	H*	H*	H*	LM	L

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT: DMMU ID: Testing Rank:	S1 H	S1-R H	S2 H	S3 H	S4 H	S5 H	US NAV	PSNS P S7 H	PROJECT S8 H	S9 H	S10 H	S11 H	S12 H	S13 H	S14 H	S15 H
METALS & ORGANOMETALS	П		П	П		П		П		П	П	П	П	П	П	
Antimony																
Arsenic Cadmium																
Copper																
Lead																
Mercury		S			S				S	S	S					S
Nickel																
Silver																ļ
Zinc																
TBT ion (porewater) LPAH																
Naphthalene																
Acenaphthene																
Acenaphthylene																
Fluorene																<u> </u>
Phenanthrene																
Anthracene 2-Methylnaphthalene																
Total LPAHs																
НРАН																
Fluoranthene																
Pyrene																
Benzo(a)anthracene																
Benzofluoranthenes (b+k) Chrysene		1					-									
Benzo(a)pyrene							 									
Indeno(1,2,3-c,d)pyrene																
Dibenzo(a,h)anthracene																
Benzo(g,h,I)perylene																
Total HPAHs																
CHLORINATED HYDROCARBONS																
1,2,4-Trichlorobenzene 1,2-Dichlorobenzene																
1,3-Dichlorobenzene																
1,4-Dichlorobenzene																
Hexaclorobenzene (HCB)																
PHTHALATES																
Bis(2-ethylheyesyl)phthalate		ļ														
Butylbenzylphthalate Di-n-butylphthalate																
Di-n-octylphthalate																
Diethylphthalate																
Dimethylphthalate																
PHENOLS																
2-Methylphenol																
4-Methylphenol 2,4-Dimethylphenol																
Pentachlorophenol																
Phenol																
MISCELANEOUS EXTRACTABLES																
Benzyl alcohol																
Benzoic acid		<u> </u>														
Dibenzofuran Hexachlorobutadiene		1					-									
Hexachlorobutadiene Hexachloroethane		1					-									
N-Nitrosodiphenylamine																-
VOLATILE ORGANICS																
Ethylbenzene																
Tetrachloroethene																
Total Zylene (sum of o,m,p) Trichloroethane																
PESTICIDES AND PCBs																
Total DDT							 									
Aldrin																
alpha-Chlordane																
Dieldrin																
Heptachlor																
gamma-BHC (Lindane) Total PCBs																
BIOASSAYS																
Amphipod																
Sediment Larval (Bivalve/Echinoderm)	2H		2H	2H	2H	2H	2H	2H	2H	2H	2H	2H	2H			2H
Neanthes Growth																
Bioassay: (Pass/Fail)		PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS
BTs eyesceeded: Bioaccumulation test conducted:	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no
Bioaccumulation (Pass/Fail):																-
ML Rule exceeded:	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no
OVERALL PASS/FAIL:		PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS
HIGHEST RANKING:	L	LM	L	L	LM	L	L	L	LM	LM	LM	L	L	L	L	LM

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

	7 S28 H
Control Contro	
Castorium Cast	
Copport Copp	
Section Sect	
Macure Section Secti	_
Saler	
Total Conversable	
TEST OF COMMING	
FAMILY	
Nagothaniene	
Accordance Acc	-
Floorence	
Pietrachisense	
Anthronome Total EPANS WEATHER TOTAL	
2.6Methyrusphralenee	
March Marc	
Figuration	-
Pyrese	
Banzo (a) partherines (P-14)	
Benzo(a) (pyrame	
Enrocalpyrene	
Benzo(a) prysee	-
Indemot 2.2 - Collywrene	-
Debrazo(gh.)Perignee	_
Total FAPA'S CLICRIMATE HYDROCARBONS 1,2.4-Incidirocherane 1,3-Dichiorobenzene 1,3-D	
CHLORNATED HYDROCARBONS	
12.2-Pichlorobenzene	
1,2-Dichlorobenzene	
1-3-Dichlorobenzene	-
1.4-Dichirobenzene (HCB)	_
BISC-2ethy-levesylphthalate	
Bisi2-etryl-leyesylphthalate Di-n-obylphthalate Di-	
Buybenzylphthalate Di-n-buylphthalate Di-n-buylphthalate Di-n-octylphthalate Di-n-octy	
Din-bryophthalate	
Dish-octypithhalate Disherhyphthalate Di	
Diethylphthalate Dimethylphthalate Dimet	
PHENOLS	
2-Methylphenol	
### ### ##############################	
2.4-Dimethylphenol S (UD) Pentachlorophenol Dehenol MISCELANEOUS EXTRACTABLES S (UD) Benzyal schol S (UD) Benzolic acid Dibenzoluran Dibenzoluran S (UD) Hexachlorobutadiene S (UD) Hexachlorobutadiene S (UD) N-Nitrosodiphenylamine S (UD) VOLATILE ORGANICS S (UD) Ethylbenzene S (UD) Intertachloroethene Total Zylene (sum of o.m.p.) Trichloroethane Total DDT Aldrin Aldrin Alpha-Chlordane Intertachlordane Dieldrin Intertachlordane BIOASSAYS	
Pentachlorophenol Phenol	-
Phenol	_
Benzyl alcohol S (UD) S (U) Benzoic acid Dibenzofuran S (UD) S (U) S (UD) S (UD	
Benzoic acid Dibenzofuran Dibe	
Dibenzofuran	
Hexachlorobutadiene	
Hexachloroethane	-
N-Nitrosodiphenylamine	_
Ethylbenzene	
Tetrachloroethene Total Zylene (sum of o,m,p) Trichloroethane PESTICIDES AND PCBS Total DDT Aldrin alpha-Chlordane Dieldrin Heptachlor gamma-BHC (Lindane) Total PCBs BIOASSAYS Amphipod Sediment Larval (Bivalve/Echinoderm) Neanthes Growth 2H Lindin Lin	
Total Zylene (sum of o,m,p)	
Trichloroethane PESTICIDES AND PCBS Total DDT	
PESTICIDES AND PCBS Total DDT Aldrin Image: Control of the cont	+
Total DDT	
Alpha-Chlordane	
Dieldrin Heptachlor	
Heptachlor	
gamma-BHC (Lindane) Total PCBs BIOASSAYS BIOASSAYS Amphipod 2H Sediment Larval (Bivalve/Echinoderm) 2H 2H Neanthes Growth 2H 2H	-
Total PCBs	+
BIOASSAYS	_
Sediment Larval (Bivalve/Echinoderm) 2H 2H 2H 2H Neanthes Growth 2H 2H Image: Control of the contr	
Neanthes Growth 2H	
	-
Bioassay: (Pass/Fail) PASS PASS PASS PASS PASS PASS PASS PAS	SS PASS
BTs eyesceded: no	
Bioaccumulation test conducted:	
Bioaccumulation (Pass/Fail):	
ML Rule exceeded: no	
HIGHEST RANKING: LM L L L L L L L L L L L L L L L L L L	

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT: DMMU ID: Testing Rank:	S29 H	S30 H	S31 H	S32 H	S33 H	S34 H	S35 H	US NAV S36 H	Y PSNS P S37 H	ROJECT S38 H	S39 H	S40 H	S41 H	S42 H	S43 H	S44 H
METALS & ORGANOMETALS			П	П	п	П	П				П			_ <u> </u>	П	П
Antimony																
Arsenic																
Cadmium																
Copper																
Lead																
Mercury																S
Nickel																
Silver																
Zinc																
TBT ion (porewater)																
LPAH																
Naphthalene																
Acenaphthene																-
Acenaphthylene																-
Fluorene Phenanthrene																-
Anthracene																+
2-Methylnaphthalene																\vdash
Total LPAHs	-							 	 	 			 	 		+-+
HPAH		1											1	1		+-+
Fluoranthene																\vdash
Pyrene																
Benzo(a)anthracene																\vdash
Benzofluoranthenes (b+k)								İ	İ	İ			İ	İ		
Chrysene								İ	İ	İ			İ	İ		
Benzo(a)pyrene																
Indeno(1,2,3-c,d)pyrene																
Dibenzo(a,h)anthracene																
Benzo(g,h,l)perylene																
Total HPAHs																
CHLORINATED HYDROCARBONS																
1,2,4-Trichlorobenzene																
1,2-Dichlorobenzene																
1,3-Dichlorobenzene																
1,4-Dichlorobenzene																
Hexaclorobenzene (HCB)																
PHTHALATES																
Bis(2-ethylheyesyl)phthalate																
Butylbenzylphthalate																
Di-n-butylphthalate																
Di-n-octylphthalate																
Diethylphthalate Dimethylphthalate																-
PHENOLS																1
2-Methylphenol																
4-Methylphenol																
2,4-Dimethylphenol																
Pentachlorophenol																
Phenol																
MISCELANEOUS EXTRACTABLES																
Benzyl alcohol																
Benzoic acid																
Dibenzofuran																
Hexachlorobutadiene																
Hexachloroethane																
N-Nitrosodiphenylamine																
VOLATILE ORGANICS																
Ethylbenzene																
Tetrachloroethene																
Total Zylene (sum of o,m,p)																ļ
Trichloroethane																Д.
PESTICIDES AND PCBs		ļ														
Total DDT								 	 	 						├
Aldrin	-	-						-	-	-						├
alpha-Chlordane Dieldrin		-						 	-	-						\vdash
Heptachlor		-						 	-	-						\vdash
gamma-BHC (Lindane)								 	 	 						++
Total PCBs								 	 	 						\vdash
BIOASSAYS								 	 	 						\vdash
Amphipod		1														
	2H				2H	2H	2H	2H	2H	2H	2H	2H	2H	2H	2H	
	2H	2H	2H													
Sediment Larval (Bivalve/Echinoderm) Neanthes Growth	2H	2H	2H		211			2H								
Sediment Larval (Bivalve/Echinoderm)	2H PASS	2H PASS	2H PASS	PASS	PASS	PASS	PASS	2H FAIL	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS
Sediment Larval (Bivalve/Echinoderm) Neanthes Growth				PASS no		PASS no	PASS no		PASS no	PASS no						PASS no
Sediment Larval (Bivalve/Echinoderm) Neanthes Growth Bioassay: (Pass/Fail) BTs eyesceeded: Bioaccumulation test conducted:	PASS	PASS	PASS		PASS			FAIL			PASS	PASS	PASS	PASS	PASS	
Sediment Larval (Bivalve/Echinoderm) Neanthes Growth Bioassay: (Pass/Fail) BTs eyesceeded: Bioaccumulation test conducted: Bioaccumulation (Pass/Fail):	PASS no	PASS no	PASS no	no	PASS no	no	no	FAIL no	no	no	PASS no	PASS no	PASS no	PASS no	PASS no	no
Sediment Larval (Bivalve/Echinoderm) Neanthes Growth Bioassay: (Pass/Fail) BTs eyesceeded: Bioaccumulation test conducted: Bioaccumulation (Pass/Fail): ML Rule exceeded:	PASS no	PASS no no	PASS no	no	PASS no	no	no	no no	no	no	PASS no	PASS no	PASS no no	PASS no no	PASS no	no
Sediment Larval (Bivalve/Echinoderm) Neanthes Growth Bioassay: (Pass/Fail) BTs eyesceeded: Bioaccumulation test conducted: Bioaccumulation (Pass/Fail):	PASS no	PASS no	PASS no	no	PASS no	no	no	FAIL no	no	no	PASS no	PASS no	PASS no	PASS no	PASS no	no

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT: DMMU ID: Testing Rank:	S45 H	S46 H	S47 H	S48 H	S48-R H	US NAV S49 H	/Y PSNS S50 H	S PROJECT S51 H	S52 H	S53 H	S54 H	S55 H	S56 H	S56-R H	S57 H	S58 H
METALS & ORGANOMETALS	п		П	П	П			П		П	П	П			П	
Antimony																
Arsenic							igwdown									
Cadmium Copper							\vdash									-
Lead									S							
Mercury	S							SB	S		S	S				
Nickel																
Silver								SB								
Zinc								S								
TBT ion (porewater)					igsquare		igsquare									igwdown
LPAH																
Naphthalene Acenaphthene					-		\vdash									-
Acenaphthylene																
Fluorene																
Phenanthrene																
Anthracene																
2-Methylnaphthalene																
Total LPAHs					ш		igspace									$ldsymbol{ldsymbol{\sqcup}}$
HPAH					igwdown		igspace								 	igspace
Fluoranthene		-		-	\longmapsto		$\vdash \vdash$	S							-	$\vdash \vdash$
Pyrene Benzo(a)anthracene					\vdash		\vdash	S							-	\vdash
Benzo(a)anthracene Benzofluoranthenes (b+k)		 			$\vdash \vdash$		$\vdash \vdash$									$\vdash \vdash$
Chrysene		1			$\vdash \vdash \vdash$		\vdash									\vdash
Benzo(a)pyrene																
Indeno(1,2,3-c,d)pyrene																
Dibenzo(a,h)anthracene																
Benzo(g,h,l)perylene																
Total HPAHs																
CHLORINATED HYDROCARBONS							ldot									ldot
1,2,4-Trichlorobenzene					igsquare			S (UD)								
1,2-Dichlorobenzene					\vdash		igwdot									igwdot
1,3-Dichlorobenzene					$\vdash \vdash$		igwdot									igwdot
1,4-Dichlorobenzene Hexaclorobenzene (HCB)					$\vdash \vdash \vdash$		$\vdash \vdash \vdash$	S (UD)						S (UD)		$\vdash \vdash \vdash$
PHTHALATES								3 (00)						3 (00)		
Bis(2-ethylheyesyl)phthalate																
Butylbenzylphthalate																
Di-n-butylphthalate																
Di-n-octylphthalate																
Diethylphthalate																
Dimethylphthalate					igsquare		igsquare									igsquare
PHENOLS 2 Mathylabasel					$\vdash \vdash$		igwdot	C (LID)								igwdot
2-Methylphenol 4-Methylphenol					-		\vdash	S (UD)								\vdash
2,4-Dimethylphenol								S (UD)								
Pentachlorophenol					\vdash			SB (UD)								
Phenol								()								
MISCELANEOUS EXTRACTABLES																
Benzyl alcohol								S (UD)								
Benzoic acid								BM								
Dibenzofuran							$ldsymbol{ldsymbol{ldsymbol{eta}}}$									$ldsymbol{ldsymbol{ldsymbol{eta}}}$
Hexachlorobutadiene								S (UD)						S (UD)		
Hexachloroethane		1		-	$\vdash \vdash$		$\vdash \vdash \vdash$	0 (110)							-	$\vdash \vdash \vdash$
N-Nitrosodiphenylamine VOLATILE ORGANICS					$\vdash \vdash \vdash$		$\vdash \vdash$	S (UD)								$\vdash \vdash$
Ethylbenzene		 			$\vdash \vdash$		$\vdash \vdash$									$\vdash \vdash$
Tetrachloroethene																
Total Zylene (sum of o,m,p)																
Trichloroethane																
PESTICIDES AND PCBs																
Total DDT					$ldsymbol{f eta}$		\Box	BM				BM				\Box
Aldrin					igspace		igspace									igspace
alpha-Chlordane		<u> </u>			igspace		igspace	S								igspace
Dieldrin				-	\longmapsto		$\vdash \vdash$	S							-	$\vdash \vdash$
Heptachlor gamma-BHC (Lindane)							\vdash								-	\vdash
Total PCBs		1			$\vdash \vdash \vdash$		$\vdash \vdash \vdash$	S		S		S			-	$\vdash \vdash \vdash$
BIOASSAYS				 	$\vdash \vdash$		$\vdash \vdash$	J		J		-			 	$\vdash \vdash$
Amphipod						ļ	2H								2H	
Sediment Larval (Bivalve/Echinoderm)		2H		2H				2H	1H	2H	2H	2H	2H		2H	
Neanthes Growth																
Bioassay: (Pass/Fail)	PASS	PASS	PASS		PASS	PASS	PASS	PASS	FAIL	PASS	PASS	PASS		PASS	FAIL	PASS
BTs eyesceeded:	no	no	no	no	no	no	no	yes	no	no	no	no	no	no	no	no
Bioaccumulation test conducted:		 			1			yes FAIL								
		1	1	I	1			FAIL	i	1		ı		1	l	
Bioaccumulation (Pass/Fail):	no	po.	po.	po.	no.	nc	no.	VAC	po.	no.	no.	no.	no.	no.	po.	no.
Bioaccumulation (Pass/Fail): ML Rule exceeded: OVERALL PASS/FAIL:	no PASS	no PASS	no PASS	no	no PASS	no PASS	no PASS	yes FAIL	no FAIL	no PASS	no PASS	no FAIL(C)	no	no PASS	no FAIL	no PASS

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT: DMMU ID: Testing Rank:	S59 H	S60 H	S61 H	S62 H	S62-R H	S63	US NAV	Y PSNS P S65 H	ROJECT S66 H	S67 H	S68 H	S69 H	S70 H	S71 H	S71-R H
METALS & ORGANOMETALS	П	П		П		П	П		П	п	П	П	П	П	П
Antimony															
Arsenic															
Cadmium Copper															-
Lead															
Mercury	S	S	S			S	S		S	S	S	S	S		
Nickel															
Silver															
Zinc															
TBT ion (porewater) LPAH			SB (E)												
Naphthalene															
Acenaphthene															
Acenaphthylene															
Fluorene															
Phenanthrene															
Anthracene															
2-Methylnaphthalene															
Total LPAHs HPAH															\vdash
Fluoranthene								 							\vdash
Pyrene															\vdash
Benzo(a)anthracene															
Benzofluoranthenes (b+k)															
Chrysene															$ldsymbol{\square}$
Benzo(a)pyrene								ļ							\vdash
Indeno(1,2,3-c,d)pyrene			-					-							\vdash
Dibenzo(a,h)anthracene Benzo(g,h,l)perylene															
Total HPAHs															-
CHLORINATED HYDROCARBONS															
1,2,4-Trichlorobenzene															
1,2-Dichlorobenzene															
1,3-Dichlorobenzene															
1,4-Dichlorobenzene															
Hexaclorobenzene (HCB)												S (UD)			.——
PHTHALATES Bis(2-ethylheyesyl)phthalate													SB (D)		
Butylbenzylphthalate													3B (D)		
Di-n-butylphthalate															
Di-n-octylphthalate															
Diethylphthalate															
Dimethylphthalate															
PHENOLS															
2-Methylphenol															.——
4-Methylphenol 2,4-Dimethylphenol															
Pentachlorophenol															_
Phenol															
MISCELANEOUS EXTRACTABLES															
Benzyl alcohol															
Benzoic acid															
Dibenzofuran															
Hexachlorobutadiene Hexachloroethane															
N-Nitrosodiphenylamine															\vdash
VOLATILE ORGANICS															
Ethylbenzene															
Tetrachloroethene															
Total Zylene (sum of o,m,p)															
Trichloroethane															<u> </u>
PESTICIDES AND PCBs	-							-							₩
Total DDT Aldrin								-							\vdash
alpha-Chlordane								 							\vdash
Dieldrin															\vdash
Heptachlor															
gamma-BHC (Lindane)															
Total PCBs						S	S			S	S		S		
BIOASSAYS		0								0					
Amphipod Sediment Langal (Rivalve/Echipoderm)	2LI	2H		211		OL.	OH.	OL!	OU.	2H	OH.	2LI		OL.	\vdash
Sediment Larval (Bivalve/Echinoderm) Neanthes Growth	2H	2H		2H		2H	2H	2H 2H	2H	2H	2H	2H		2H	
Bioassay: (Pass/Fail)	PASS	FAIL	PASS		PASS	PASS	PASS	FAIL	PASS	FAIL	PASS	PASS	PASS	PASS	PASS
BTs eyesceeded:	no	no	no	no	no	no	no	no	no	no	no	no	yes	no	no
Bioaccumulation test conducted:													no		
Bioaccumulation (Pass/Fail):															igspace
ML Rule exceeded:	no	no	no EAIL (C)	no	no DACC	no DACC	no DACC	no	no DACC	no	no DACC	no DACC	no FAIL(C)	no	no DACC
OVERALL PASS/FAIL: HIGHEST RANKING:	PASS LM	FAIL H*	FAIL(C) H	L	PASS	PASS LM	PASS LM	FAIL H*	PASS LM	FAIL H*	PASS LM	PASS LM	FAIL(C)	1	PASS
INGILOT KANKING.	LIVI	rī	п	L	L	LIVI	LIVI	п	LIVI	H*	LIVI	LIVI	п	L	L

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT: DMMU ID: Testing Rank:	S72 H	S73 H	S73-R H	S77 H	S78 H	\$80 H	US NAVY D1 H	/ PSNS P D2 H	ROJECT D3 H	D3-R H	D4a H	D4b H	D4c H	D4d H	D5 H	D6a H
METALS & ORGANOMETALS		П		П		П		П			п			П		
Antimony																
Arsenic				S												
Cadmium																
Copper				S	S											
Lead Mercury	S			ВМ	BM		S									
Nickel	3			DIVI	DIVI		3									
Silver																
Zinc					S											
TBT ion (porewater)																
LPAH																
Naphthalene																
Acenaphthene	S															
Acenaphthylene																
Fluorene	S															
Phenanthrene	S (D)															
Anthracene	S				ļ											
2-Methylnaphthalene Total LPAHs		-	 		-		 								-	
HPAH		-	1		-		1		-	1			1		-	
Fluoranthene	SB (D)				1				 	 			 			
Pyrene	S (D)				-											
Benzo(a)anthracene	S (D)		1		1		1									
Benzofluoranthenes (b+k)	- (-)									l			l			
Chrysene	S (D)									l			l			
Benzo(a)pyrene																
Indeno(1,2,3-c,d)pyrene	S															
Dibenzo(a,h)anthracene																
Benzo(g,h,l)perylene																
Total HPAHs	S															
CHLORINATED HYDROCARBONS																
1,2,4-Trichlorobenzene																
1,2-Dichlorobenzene																
1,3-Dichlorobenzene																
1,4-Dichlorobenzene																
Hexaclorobenzene (HCB)				S (UD)	S (UD)											
PHTHALATES					ļ											
Bis(2-ethylheyesyl)phthalate																
Butylbenzylphthalate Di-n-butylphthalate					1											
Di-n-octylphthalate																
Diethylphthalate																
Dimethylphthalate																
PHENOLS																
2-Methylphenol																
4-Methylphenol																
2,4-Dimethylphenol				S (UD)	S (UD)											
Pentachlorophenol																
Phenol																
MISCELANEOUS EXTRACTABLES																
Benzyl alcohol				S (UD)												
Benzoic acid										 			 			
Dibenzofuran				0.015	0 "					 			 			
Hexachloropthana		-		S (UD)	S (UD)				 	 			 			
Hexachloroethane N-Nitrosodiphenylamine		-	-	S (UD)	S (UD)		-		 	 			 		-	
VOLATILE ORGANICS		 		3 (00)	3 (UD)				 	1			1			
Ethylbenzene		-			-											
Tetrachloroethene										l			l			
Total Zylene (sum of o,m,p)										l			l			
Trichloroethane			1				1									
PESTICIDES AND PCBs																
Total DDT																
Aldrin																
alpha-Chlordane																
Dieldrin																
Heptachlor						-										
gamma-BHC (Lindane)										ļ			ļ			
Total PCBs	S			S	S					ļ			ļ			
BIOASSAYS										1			1			
Amphipod	1H			2H	1H		2H	0	011	<u> </u>					1H	
Sediment Larval (Bivalve/Echinoderm)	2H			1H	1H	2H	1H	2H	2H	 			2H	2H	2H	
Neanthes Growth Bioassay: (Pass/Fail)	FAIL		PASS	2H FAIL	FAIL	PASS	FAIL	PASS	-	PASS	PASS	PASS	PASS	PASS	2H FAIL	PASS
BTs eyesceeded:	yes	no	no	yes	yes	no	no	no no	no	no	no	no	no	no	no	no
Bioaccumulation test conducted:	no	110	110	no	no	110	110	110	110	110	110	110	110	110	110	110
Bioaccumulation (Pass/Fail):	110			110	110					1			1			
ML Rule exceeded:	no	no	no	yes	yes	no	no	no	no	no	no	no	no	no	no	no
OVERALL PASS/FAIL:	FAIL		1	FAIL	FAIL	PASS	FAIL	PASS		PASS	PASS	PASS	PASS	PASS	FAIL	PASS

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT: DMMU ID: Testing Rank:	D6b H	D6c H	US NAV D8 H	/Y PSNS SS101 H	PROJEC SS102 H	T SS103 H	S2B H	S3 H	JAMES H S5 H	IARDIE G' CB H	YPSUM CC H	CD H	CE H	OLYMPIA B1 H	A HARB TBW1 H
METALS & ORGANOMETALS															
Antimony															
Arsenic															
Cadmium															
Copper															
Lead												S			
Mercury Nickel												3			
Silver															
Zinc															
TBT ion (porewater)														SB	SB
LPAH															
Naphthalene															
Acenaphthene															
Acenaphthylene															
Fluorene															
Phenanthrene											S				
Anthracene															
2-Methylnaphthalene	-				 			-					 	-	
Total LPAHs HPAH		-			 			 				-	 		-
Fluoranthene	-	1			-			-	1		S		-	1	1
Pyrene					 			 			S		 		
Benzo(a)anthracene		1			1			1	1				1	1	1
Benzofluoranthenes (b+k)															
Chrysene															
Benzo(a)pyrene															
Indeno(1,2,3-c,d)pyrene															
Dibenzo(a,h)anthracene															
Benzo(g,h,l)perylene															
Total HPAHs															
CHLORINATED HYDROCARBONS															
1,2,4-Trichlorobenzene															
1,2-Dichlorobenzene															
1,3-Dichlorobenzene															
1,4-Dichlorobenzene															
Hexaclorobenzene (HCB)															
PHTHALATES															
Bis(2-ethylheyesyl)phthalate															
Butylbenzylphthalate Di-n-butylphthalate															
Di-n-octylphthalate															
Diethylphthalate															
Dimethylphthalate															
PHENOLS															
2-Methylphenol															
4-Methylphenol															
2,4-Dimethylphenol															
Pentachlorophenol															
Phenol															
MISCELANEOUS EXTRACTABLES															
Benzyl alcohol															
Benzoic acid															
Dibenzofuran													ļ		
Hexachlorobutadiene Hexachloroethane	-				 			-					 	-	
N-Nitrosodiphenylamine		-			 			 				<u> </u>	 		-
VOLATILE ORGANICS	H				 										
Ethylbenzene															
Tetrachloroethene															
Total Zylene (sum of o,m,p)															
Trichloroethane		İ			İ			İ				İ	İ	1	İ
PESTICIDES AND PCBs															
Total DDT															
Aldrin															
alpha-Chlordane															
Dieldrin												S			
Heptachlor															
gamma-BHC (Lindane)															
Total PCBs					 		S	S	S	S	S	SB	SB		
BIOASSAYS		1	4.1				411	411	011	211		011			I
Amphipod Sodiment Lengt (Biyolya/Echinodorm)	011	OL I	1H	01.1	OL.	OL.	1H	1H	2H	2H	OL I	2H	 		-
Sediment Larval (Bivalve/Echinoderm) Neanthes Growth	2H	2H	2H	2H	2H	2H		-	1H		2H 2H	1H		1	1
Bioassay: (Pass/Fail)	PASS	PASS	FAIL	PASS	PASS	PASS	FAIL	FAIL	FAIL	PASS	FAIL	FAIL	PASS	NA	NA
BTs eyesceeded:	no	no	no	no	no	no	no	no	no	no	no	yes	yes	yes	yes
Bioaccumulation test conducted:	110	110	110	110	110	110	110	110	110	110	110	no	no	yes	yes
Bioaccumulation (Pass/Fail):					1			1					- ···	PASS	PASS
ML Rule exceeded:	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no
OVERALL PASS/FAIL:	PASS	PASS	FAIL	PASS	PASS	PASS	FAIL	FAIL	FAIL	PASS	FAIL	FAIL	PASS	PASS	PASS
HIGHEST RANKING:			H*						H*					LM	LM

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

### A PART	PROJECT:	S1	S2	S3	USN PIT	-CAD CH/ S5	ARACTER S6	RIZATION S7	S8	S9	S10	S11	S12	S13	S14	S15	S16
ASSESSION	Testing Rank:																Н
Agents																	
Control																	
Company																	
Marcuny																	
Section																	
State		S	S				S	S		S	S	S	S				S
Table																	\vdash
Till Care processors																	
Registrations																	
Accordation																	
According																	—
Process																	
Martinerine																	
Extension programme																	
Trots (PAPS)																	
Figuration																	
Flacenombres		l .	1							1							
Provide Prov																	
Bearcolloyprene	Pyrene																
Empress	Benzo(a)anthracene																
Bezond Section Secti		 	-		ļ											ļ	
Indicated 12-28-city/princeme																	
Distraction Alperformance		l								1							
Total IP/Ai-Nis CML CRIMATED HYDROCARBONS 1, 2-f. Trinifinoteduruses 1, 2-f																	
Committee Comm	Benzo(g,h,l)perylene																
12.4-Trichrobotescee																	
1.2-Dichlorobenzene																	-
1.5-Dichiorobenzene																	\vdash
Hexandrotename (HCES)																	
PITHALATES																	
Bis/Jebray/phrabiate																	
Butybensylphthalates																	
Dentrophythalate																	
Distript/Inhalate																	
Dimetrylyphenoid																	
PHENOLS																	—
AMethylphenol		-															\vdash
### Admity/phenol																	
Pentachlorophenol																	
Phenol																	
Benzyl alcohol Benz			ļ														—
Benzyl alcohol Benzol acid																	\vdash
Dibenzoturan																	
Hexachlorobutadiene																	
Hexachloroethane																	
N-Nitrosodiphenylamine		 	1							 							\vdash
VOLATILE ORGANICS		 								<u> </u>							\vdash
Ethylbenzene	VOLATILE ORGANICS																
Total Zylene (sum of o,m,p) Trichlorethane	Ethylbenzene																
Trichloroethane			1														
PESTICIDES AND PCBs		 	-							 							\vdash
Total DDT		 	1							 							
Aldrin alpha-Chlordane Dieldrin Heptachlor gamma-BHC (Lindane) Total PCBs BIOASSAYS Amphipod Sediment Larval (Bivalve/Echinoderm) PASS PASS FAIL PASS PASS PASS PASS PASS PASS PASS PAS		l															
Dieldrin Heptachior Gamma-BHC (Lindane)	Aldrin																
Heptachlor gamma-BHC (Lindane)																	
Gamma-BHC (Lindane) Total PCBs September Total PCBs September Total PCBs September Total PCBs September Total PCBs September Total PCBs September		-	1		-					-						-	\vdash
Total PCBs BIOASSAYS		 								 							\vdash
BIOASSAYS		l								1							
Sediment Larval (Bivalve/Echinoderm)		İ															\Box
Neanthes Growth 2H																	
Bioassay: (Pass/Fail) PASS		2H	2H		2H	2H		2H	2H		2H			2H	2H		2H
BTs eyesceeded:		PASS	PASS		PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS.	DASS	PASS	PASS	PASS	PASS
Bioaccumulation test conducted:																	no
Bioaccumulation (Pass/Fail):		110	110	110	110	110	110	110	110	110	110	110	110	110	110	110	110
OVERALL PASS/FAIL: PASS PASS FAIL PASS PASS PASS PASS PASS PASS PASS PAS	Bioaccumulation (Pass/Fail):																
																	no
IN I LW I LW I LW I TO I L I LM I LW I LM I LM I LM I LM I LM																	PASS
	HIGHEST KANKING:	LIVI	LM	П.			LIVI	LIVI		LIVI	LIVI	LIVI	LIVI	L			LIVI

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT: DMMU ID: Testing Rank:	S17 H	S18 H	S19 H	S20 H	USN PIT- S21 H	-CAD CH/ S22 H	ARACTER S23 H	S24 H	S25 H	S26 H	S27 H	S28 H	S29 H	S30 H	DUWAMI S1 H	ISH O&M S3 H
METALS & ORGANOMETALS																
Antimony																
Arsenic																
Cadmium																
Copper																-
Lead Mercury	S	S				S	S	S		S	S		S	S		
Nickel	3	3				3	3	3		3	3		3	3		
Silver																
Zinc																
TBT ion (porewater)																SB
LPAH																
Naphthalene																l
Acenaphthene																
Acenaphthylene																
Fluorene																
Phenanthrene																
Anthracene 2-Methylnaphthalene																
Total LPAHs																
HPAH																ſ
Fluoranthene																ſ
Pyrene																
Benzo(a)anthracene																
Benzofluoranthenes (b+k)																
Chrysene																
Benzo(a)pyrene																<u> </u>
Indeno(1,2,3-c,d)pyrene							ļ									—
Dibenzo(a,h)anthracene				-			-			-		-	-			
Benzo(g,h,l)perylene																
Total HPAHs CHLORINATED HYDROCARBONS																-
1,2,4-Trichlorobenzene																-
1,2-Dichlorobenzene																
1,3-Dichlorobenzene																
1,4-Dichlorobenzene																
Hexaclorobenzene (HCB)																
PHTHALATES																
Bis(2-ethylheyesyl)phthalate																
Butylbenzylphthalate																
Di-n-butylphthalate																
Di-n-octylphthalate Diethylphthalate																-
Dimethylphthalate																
PHENOLS																
2-Methylphenol																
4-Methylphenol																
2,4-Dimethylphenol																
Pentachlorophenol																
Phenol																l
MISCELANEOUS EXTRACTABLES																
Benzyl alcohol																
Benzoic acid							 							-	—	
Dibenzofuran Hexachlorobutadiene	-	1		-			-			-		-	-	-		
Hexachloroethane	\vdash	 		 			 			 		-	 	 	\vdash	
N-Nitrosodiphenylamine																
VOLATILE ORGANICS																
Ethylbenzene																ſ
Tetrachloroethene																
Total Zylene (sum of o,m,p)																
Trichloroethane																
PESTICIDES AND PCBs																
Total DDT																
Aldrin							ļ									
alpha-Chlordane							ļ									—
Dieldrin	ļ	-		-						-			-			
Heptachlor gamma-BHC (Lindane)				-			 			-		-	-	 		—
Total PCBs	-	1		-			-			-		-	-	-	S	
BIOASSAYS															3	
Amphipod																ł
Sediment Larval (Bivalve/Echinoderm)	2H	2H	2H	2H		2H	2H	2H	2H	2H	2H	2H	2H	2H	2H	
Neanthes Growth															2H	
Bioassay: (Pass/Fail)	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	FAIL	NA
BTs eyesceeded:	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no	yes
Bioaccumulation test conducted:																no
Bioaccumulation (Pass/Fail):																<u> </u>
ML Rule exceeded:	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no
OVERALL PASS/FAIL:	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	PASS	FAIL	FAIL(C)
HIGHEST RANKING:	LM	LM	L	L	L	LM	LM	LM	L	LM	LM	L	LM	LM	H*	H*

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT: DMMU ID: Testing Rank:	S4 H	S5 H	S6 H	S7 H	DUWAMI S8 H	ISH O&M S9 H	S10 H	S11 H	S12 H	S13 H	S14 H	S15 H	S16 H	B1 H	B2 H	WEYER B1 H
METALS & ORGANOMETALS																
Antimony		† ·						·		·		1				
Arsenic																
Cadmium		<u> </u>		-	 	 	ļ!	 		 						
Copper		<u> </u>		-		├ ──┤	-	 	 	 						1
Lead Mercury																
Nickel																
Silver																
Zinc																
TBT ion (porewater)	SB (M)											SB				SB
LPAH																
Naphthalene		<u> </u>					 	 	 	 						_
Acenaphthene Acenaphthylene					\vdash	\vdash		—	 	—						1
Fluorene		-							 			-		-		
Phenanthrene																
Anthracene																
2-Methylnaphthalene																
Total LPAHs		<u> </u>														
HPAH		<u> </u>	igsquare	 	igsquare							 		<u> </u>		<u> </u>
Fluoranthene	-	 	$\vdash \vdash \vdash$	 	$\vdash \vdash \vdash$	$\vdash \vdash \vdash$	$\vdash \vdash \vdash$	 	 	 			 	—	 	
Pyrene Benzo(a)anthracene	-	 	\vdash	 	$\vdash \vdash \vdash$	$\vdash \vdash$	\vdash				 	 	\vdash	 	\vdash	
Benzofluoranthenes (b+k)		 	\vdash						\vdash		 			+		S
Chrysene																S
Benzo(a)pyrene																
Indeno(1,2,3-c,d)pyrene																
Dibenzo(a,h)anthracene																
Benzo(g,h,I)perylene		<u> </u>														
Total HPAHs		ļ!			<u> </u>	igwdapsilon										
CHLORINATED HYDROCARBONS					igwdown	\vdash	-	 	 	 						
1,2,4-Trichlorobenzene 1,2-Dichlorobenzene		 			\vdash	\vdash		— —		— —						-
1,3-Dichlorobenzene		-							 			-		-		_
1,4-Dichlorobenzene																
Hexaclorobenzene (HCB)																S (U)
PHTHALATES																
Bis(2-ethylheyesyl)phthalate																
Butylbenzylphthalate																
Di-n-butylphthalate		<u> </u>		-	 	 										
Di-n-octylphthalate		<u> </u>					 	 	 	 						
Diethylphthalate Dimethylphthalate																-
PHENOLS																
2-Methylphenol																
4-Methylphenol																
2,4-Dimethylphenol																
Pentachlorophenol																
Phenol																
MISCELANEOUS EXTRACTABLES					igwdown	\vdash	-	 	 	 						
Benzyl alcohol		<u> </u>					 	 	 	 						_
Benzoic acid Dibenzofuran	-	\vdash	$\vdash \vdash$	 	$\vdash \vdash$	$\vdash \vdash$	\vdash		\vdash			\vdash	$\vdash \vdash$	\vdash	\vdash	\vdash
Hexachlorobutadiene		 	\vdash						\vdash		 			+		S (U)
Hexachloroethane								[[<u> </u>		J (J)
N-Nitrosodiphenylamine																S (J)
VOLATILE ORGANICS																
Ethylbenzene						\Box										
Tetrachloroethene		<u> </u>	igsquare		igsquare	 _		 	لـــــــــا	 		 	<u> </u>	<u> </u>	<u> </u>	
Total Zylene (sum of o,m,p)	-	 	$\vdash \vdash \vdash$	 	$\vdash \vdash \vdash$	$\vdash \vdash \vdash$	$\vdash \vdash \vdash$	 	├	 			 	-	 	├
Trichloroethane PESTICIDES AND PCBs		├──		-	$\vdash \vdash$	$\vdash \vdash \vdash$			\vdash		—	 	 	<u> </u>	 	+-+
Total DDT		\vdash	$\vdash \vdash$	 	$\vdash \vdash$		$\vdash \vdash$		\vdash		 	\vdash	$\vdash \vdash$	\vdash	$\vdash \vdash$	S (U)
Aldrin		 	\vdash		\vdash	\vdash			\vdash		+			+		3 (0)
alpha-Chlordane																\vdash
Dieldrin																
Heptachlor																
gamma-BHC (Lindane)																
Total PCBs		<u> </u>	S	S	S	S		S	S	S	S	S	S	S	S	S
BIOASSAYS]				i l		i '		i ']		
Amphipod	-	 	$\vdash \vdash \vdash$	-	$\vdash \vdash$	$\vdash \vdash$	 	 	011	 			<u> </u>	<u> </u>	011	2H
Sediment Larval (Bivalve/Echinoderm) Neanthes Growth	-	211		211	211		$\vdash \vdash \vdash$	211	2H		<u> </u>	 		211	2H	2H
Bioassay: (Pass/Fail)	NA	2H PASS	PASS	2H PASS	2H PASS	PASS	PASS	2H PASS	PASS	PASS	PASS	NA	PASS	2H PASS	2H FAIL	FAIL
BTs eyesceeded:	yes	no	no	no	no	no	no	no	no	no	no	yes	no	no	no	yes
Bioaccumulation test conducted:	no	110	110	110	110	110	110	110	110	110	110	no	110	110	110	no
Bioaccumulation (Pass/Fail):																
ML Rule exceeded:	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no
OVERALL PASS/FAIL:		2400	D400	PASS	PASS	PASS	DACC	PASS	PASS	PASS	PASS	FAIL(C)	PASS	DACC	FAIL	FAIL
HIGHEST RANKING:	FAIL(C) H*	PASS L	PASS LM	LM	LM	LM	PASS L	LM	LM	LM	LM	H*	LM	PASS LM	H*	H*

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT:		WEYERI	HAEUSER	R CO. HW	DG						MANKE	LUMBER	HWDG			
DMMU ID	B2	B3	B5	B6	B7	В8	В9	B10	A1	A2	A3	A4	A5	A6	A7	A8
Testing Rank	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н
METALS & ORGANOMETALS																
Antimony Arsenic									S	S	S	S		S		
Cadmium																
Copper																
Lead																
Mercury														S		
Nickel Silver																
Zinc									S			S				
TBT ion (porewater)	SB		SB	SB	SB	SB										
LPAH																
Naphthalene																
Acenaphthene								S							S	
Acenaphthylene Fluorene															S	
Phenanthrene								S							S	
Anthracene								_							S	
2-Methylnaphthalene																
Total LPAHs								S							S	
HPAH								S	S		S		CD		CD	
Fluoranthene Pyrene		 		 				S	3		3		SB S		SB S	S
Benzo(a)anthracene								S					S		S	
Benzofluoranthenes (b+k)	S					S							S		S	
Chrysene	S			S		S	S	S	S		S		S		S	
Benzo(a)pyrene																
Indeno(1,2,3-c,d)pyrene	S		_	<u> </u>				S	1					-	-	0.// 0.
Dibenzo(a,h)anthracene Benzo(g,h,l)perylene	-		S						-					-		S (U)
Total HPAHs													S		S	
CHLORINATED HYDROCARBONS		1							İ							
1,2,4-Trichlorobenzene																
1,2-Dichlorobenzene																
1,3-Dichlorobenzene																
1,4-Dichlorobenzene	0.40	0.40	0.40	0.40			0 4 11	0.00		0.40	0.40	0.40	0.40	0.40	0.40	D14 (1)
Hexaclorobenzene (HCB) PHTHALATES	S (U)	S (U)	S (U)	S (U)			S (U)	S (U)		S (U)	S (U)	S (U)	S (U)	S (U)	S (U)	BM (U)
Bis(2-ethylheyesyl)phthalate																
Butylbenzylphthalate																
Di-n-butylphthalate																
Di-n-octylphthalate																
Diethylphthalate																
Dimethylphthalate PHENOLS																
2-Methylphenol																
4-Methylphenol																
2,4-Dimethylphenol		S (U)	S (U)	S (U)				S (U)			S (U)					
Pentachlorophenol	SB													SB (U)		BM
Phenol																
MISCELANEOUS EXTRACTABLES	C (II)		C (LI)	S (U)			S (U)	S (U)		S (U)		S (U)	S (U)	S (U)	S (U)	C (LI)
Benzyl alcohol Benzoic acid	S (U)		S (U)	S (U)			3 (0)	S (U)		3 (0)		S (U)	3 (0)	S (U)	S (U)	S (U)
Dibenzofuran															S	
Hexachlorobutadiene	S (U)	S (U)	S (U)	S (U)			S (U)	S (U)		S (U)	S (U)	S (U)	S (U)	S (U)	S (U)	SB (U)
Hexachloroethane																
N-Nitrosodiphenylamine	S (U)	S (U)	S (U)	S (U)			S (U)	S (U)		S (U)	S (U)	S (U)	S (U)	S (U)	S (U)	BM (U)
VOLATILE ORGANICS	-	1	-	1					 					-	1	
Ethylbenzene Tetrachloroethene		 		1					 						 	
Total Zylene (sum of o,m,p)																
Trichloroethane																
PESTICIDES AND PCBs																
Total DDT	S (U)	S (J)	S (U)			S (U)	S (U)		S (U)		S (UJ)		S (U)	S (U)	S (U)	
Aldrin	3 (0)			1						 						
	3 (0)															1
alpha-Chlordane	3 (0)															
alpha-Chlordane Dieldrin	3 (0)															
alpha-Chlordane Dieldrin Heptachlor	3 (0)															
alpha-Chlordane Dieldrin	S		S			S	S		S		S		S		S	
alpha-Chlordane Dieldrin Heptachlor gamma-BHC (Lindane) Total PCBs BIOASSAYS	S		S			S	S				S				S	
alpha-Chlordane Dieldrin Heptachlor gamma-BHC (Lindane) Total PCBs BIOASSAYS Amphipod	S 1H	1H	S			1H			1H	1H		1H	1H	1H		
alpha-Chlordane Dieldrin Heptachlor gamma-BHC (Lindane) Total PCBs BIOASSAYS Amphipod Sediment Larval (Bivalve/Echinoderm)	S	1H	S				S 2H	2H		1H	S 2H	1H 2H		1H 2H	S 1H	
alpha-Chlordane Dieldrin Heptachlor gamma-BHC (Lindane) Total PCBs BIOASSAYS Amphipod Sediment Larval (Bivalve/Echinoderm) Neanthes Growth	S 1H 2H			Diece	DACC	1H 2H	2H		1H 2H		2H	2H	1H 2H	2H	1H	NA
alpha-Chlordane Dieldrin Heptachlor gamma-BHC (Lindane) Total PCBs BIOASSAYS Amphipod Sediment Larval (Bivalve/Echinoderm) Neanthes Growth Bioassay: (Pass/Fail)	S 1H 2H	FAIL	PASS	PASS	PASS	1H 2H FAIL	2H PASS	PASS	1H 2H FAIL	FAIL	2H PASS	2H FAIL	1H 2H FAIL	2H FAIL	1H FAIL	NA Ves
alpha-Chlordane Dieldrin Heptachlor gamma-BHC (Lindane) Total PCBs BIOASSAYS Amphipod Sediment Larval (Bivalve/Echinoderm) Neanthes Growth Bioassay: (Pass/Fail) BTs eyesceeded:	S 1H 2H FAIL yes		PASS yes	yes	yes	1H 2H FAIL yes	2H		1H 2H		2H	2H	1H 2H FAIL yes	2H FAIL yes	1H FAIL yes	yes
alpha-Chlordane Dieldrin Heptachlor gamma-BHC (Lindane) Total PCBs BIOASSAYS Amphipod Sediment Larval (Bivalve/Echinoderm) Neanthes Growth Bioassay: (Pass/Fail)	S 1H 2H	FAIL	PASS			1H 2H FAIL	2H PASS	PASS	1H 2H FAIL	FAIL	2H PASS	2H FAIL	1H 2H FAIL	2H FAIL	1H FAIL	
alpha-Chlordane Dieldrin Heptachlor gamma-BHC (Lindane) Total PCBs BIOASSAYS Amphipod Sediment Larval (Bivalve/Echinoderm) Neanthes Growth Bioassay: (Pass/Fail) BTs eyesceded: Bioaccumulation test conducted:	S 1H 2H FAIL yes	FAIL	PASS yes yes	yes yes	yes yes	1H 2H FAIL yes	2H PASS	PASS	1H 2H FAIL	FAIL	2H PASS	2H FAIL	1H 2H FAIL yes	2H FAIL yes	1H FAIL yes	yes
alpha-Chlordane Dieldrin Heptachlor gamma-BHC (Lindane) Total PCBs BIOASSAYS Amphipod Sediment Larval (Bivalve/Echinoderm) Neanthes Growth Bioassay: (Pass/Fail) BTs eyesceedde: Bioaccumulation test conducted: Bioaccumulation (Pass/Fail):	S 1H 2H FAIL yes no	FAIL no	PASS yes yes PASS	yes yes PASS	yes yes PASS	1H 2H FAIL yes no	2H PASS no	PASS no	1H 2H FAIL no	FAIL no	2H PASS no	2H FAIL no	1H 2H FAIL yes no	FAIL yes no	1H FAIL yes no	yes no

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT:						MANKE I	LUMBER	HWDG							
DMMU ID: Testing Rank:	A11 H	A12 H	A13 H	A14 H	A15 H	A16 H	A17 H	A18 H	A19 H	A21 H	A22 H	A23 H	A25 H	A26 H	A27 H
METALS & ORGANOMETALS			П	П	П	П	П			П	П	П		П	
Antimony															
Arsenic Cadmium												S	S		
Copper															
Lead															
Mercury															
Nickel Silver															
Zinc													S		
TBT ion (porewater)															
LPAH															
Naphthalene Acenaphthene									S						
Acenaphthylene															
Fluorene									S						
Phenanthrene Anthracene				S					S						
2-Methylnaphthalene									3						
Total LPAHs															
HPAH															
Fluoranthene Pyrene	-	}		S				}	S					S	
Benzo(a)anthracene	l														
Benzofluoranthenes (b+k)															
Chrysene	S		S	S					S						
Benzo(a)pyrene Indeno(1,2,3-c,d)pyrene	 														
Dibenzo(a,h)anthracene	l														
Benzo(g,h,l)perylene															
Total HPAHs	S			S					S						
CHLORINATED HYDROCARBONS 1,2,4-Trichlorobenzene															
1,2,4-1 richiorobenzene 1,2-Dichlorobenzene															
1,3-Dichlorobenzene															
1,4-Dichlorobenzene															
Hexaclorobenzene (HCB) PHTHALATES	S (U)	S (U)	S (U)	SB (U)	S (U)	S (U)	S (U)		S (U)					S (U)	
Bis(2-ethylheyesyl)phthalate															
Butylbenzylphthalate															
Di-n-butylphthalate															
Di-n-octylphthalate															
Diethylphthalate Dimethylphthalate															
PHENOLS															
2-Methylphenol															
4-Methylphenol 2,4-Dimethylphenol					S (U)										-
Pentachlorophenol	SB (U)			BM (U)	3 (0)				S (U)						
Phenol				\-/											
MISCELANEOUS EXTRACTABLES															
Benzyl alcohol Benzoic acid	S (U)	S (U)		S (U)	S (U)		S (U)		S (U)				S (U)		
Dibenzofuran									S (U)						
Hexachlorobutadiene	S (U)	S (U)	S (U)	S (U)	S (U)	S (U)	S (U)		S (U)				S (U)		
Hexachloroethane		0.40	0.40	DM	0.45	0.45	0.40		0.40				0.40		
N-Nitrosodiphenylamine VOLATILE ORGANICS	l	S (U)	S (U)	BM (U)	S (U)	S (U)	S (U)		S (U)				S (U)		
Ethylbenzene															
Tetrachloroethene															
Total Zylene (sum of o,m,p)	-														
Trichloroethane PESTICIDES AND PCBs	 														
Total DDT	S (U)		S (U)	S (U)	S (U)	S (U)	S (U)	S (U)	S (U)		S (UJ)				
Aldrin															
alpha-Chlordane	0 (11)														\vdash
Dieldrin Heptachlor	S (U)														
gamma-BHC (Lindane)									S (U)						
Total PCBs	S	S	S	S	S	S	S	S	S		S	S		S	S
BIOASSAYS															
Amphipod Sediment Larval (Bivalve/Echinoderm)	2H 2H	1H		1H 2H		1H 2H			2H 2H	1H 2H	1H		2H		2H
Neanthes Growth	ZH			211		201			20	201			211		2/1
Bioassay: (Pass/Fail)	FAIL	FAIL	PASS	FAIL	PASS	FAIL	PASS	PASS	FAIL	FAIL	FAIL	PASS	PASS	PASS	PASS
BTs eyesceeded:	yes	no	no	yes	no	no	no	no	no	no	no	no	no	no	no
Bioaccumulation test conducted: Bioaccumulation (Pass/Fail):	no	-		no				-							1
ML Rule exceeded:	no	no	no	yes	no	no	no	no	no	no	no	no	no	no	no
OVERALL PASS/FAIL:	FAIL	FAIL	PASS	FAIL	PASS	FAIL	PASS	PASS	FAIL	FAIL	FAIL	PASS	PASS	PASS	PASS
HIGHEST RANKING:	Н	H*	LM	Н	LM	H*	LM	LM	H*	H*	H*	LM	LM	LM	LM

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT: DMMU ID: Testing Rank:	POA-CAI C8 M	P SANTE I COMP1 M	MARINA COMP2 M	POB-PAD C1 H	C2 LM	POT-HYI C4 H	LEBOS/BL C9 LM	AIR SLIP C10A LM	ONE (MO	OUTH OI C12 LM	F HYLEE C16 LM	BOS) C17 LM	C43 H	(WAS.W) C37 H	POT-HY C18 H	'L(MP) C19 H
METALS & ORGANOMETALS																
Antimony																
Arsenic																
Cadmium																
Copper																
Lead								S		S						
Mercury Nickel								3		3						
Silver																
Zinc																
TBT ion (porewater)		SB	SB					SB (G)		SB (G)				SB (G)		SB (G)
LPAH		- OD	05					05 (0)		02 (0)				<i>CD</i> (C)		05 (0)
Naphthalene	S					M										
Acenaphthene				S		M										
Acenaphthylene																
Fluorene				S		M										
Phenanthrene				S		M										
Anthracene						S										
2-Methylnaphthalene				S		M										
Total LPAHs				S		М										
НРАН																
Fluoranthene	S			S		BM										
Pyrene	S					M										
Benzo(a)anthracene						M				ļ						
Benzofluoranthenes (b+k)						M										
Chrysene	S					S				ļ						
Benzo(a)pyrene						BM			ļ	ļ						
Indeno(1,2,3-c,d)pyrene						M	ļ		ļ	 						
Dibenzo(a,h)anthracene						S	ļ		ļ	 						
Benzo(g,h,l)perylene						M	ļ		 	 						
Total HPAHs	S															
CHLORINATED HYDROCARBONS																
1,2,4-Trichlorobenzene								S								
1,2-Dichlorobenzene																
1,3-Dichlorobenzene																
1,4-Dichlorobenzene					0.00	0.40	0.00	-	-	-		0.00				
Hexaclorobenzene (HCB)					S (M)	S (U)	S (M)	S	S	S		S (M)				
PHTHALATES																
Bis(2-ethylheyesyl)phthalate																
Butylbenzylphthalate																
Di-n-butylphthalate Di-n-octylphthalate																
Diethylphthalate																
Dimethylphthalate																
PHENOLS																
2-Methylphenol						S (U)										
4-Methylphenol						0 (0)										
2,4-Dimethylphenol						S (U)										
Pentachlorophenol						S										
Phenol																
MISCELANEOUS EXTRACTABLES																
Benzyl alcohol						S (U)										
Benzoic acid						S										
Dibenzofuran						М										
Hexachlorobutadiene					S	S (U)	S	S	S	S		S				
Hexachloroethane						(-)									1	
N-Nitrosodiphenylamine						S (U)										
VOLATILE ORGANICS																
Ethylbenzene																
Tetrachloroethene																
Total Zylene (sum of o,m,p)																
Trichloroethane																
PESTICIDES AND PCBs																
Total DDT					S		S	S (NG)	S (NG)	S (NG)	S (NG)			S	S	S
Aldrin											S (U)	S (U)				
alpha-Chlordane											S (U)	S (U)				
Dieldrin								S (Y)	S (Y)	S (Y)		S (U)			S (Y)	S (Y)
Heptachlor											S (U)	S (U)				
gamma-BHC (Lindane)								S	S	S (Y)		S				
Total PCBs						S		S (Y)	S (Y)		S (UE)	S (U)	S	S	S	S
BIOASSAYS																
Amphipod										2H	2H	2H	2H	2H		
Sediment Larval (Bivalve/Echinoderm)							1H	1H	1H	1H	1H	1H			1H	1H
Neanthes Growth								2H	2H							
Bioassay: (Pass/Fail)	PASS	NA	NA	PASS	PASS	NA	FAIL	FAIL	FAIL	FAIL	FAIL	FAIL	PASS	PASS	FAIL	FAIL
BTs eyesceeded:	no	yes	yes	no	no	yes	no	yes	no	yes	no	no	no	yes	no	yes
							1	no		no				no		no
Bioaccumulation test conducted:		yes	yes			no		110		110				110		
Bioaccumulation (Pass/Fail):		yes PASS	yes PASS			no		110		110				110		
Bioaccumulation (Pass/Fail): ML Rule exceeded:	no			no	no	yes	no	no	no	no	no	no	no	no	no	no
Bioaccumulation (Pass/Fail):		PASS	PASS	no PASS LM	no PASS LM		no FAIL H		no FAIL H		no FAIL H	no FAIL H	no PASS LM		no FAIL H*	no FAIL H*

APPENDIX C - DY 00/01 EVALUATION GUIDELINE EXCEEDANCES

PROJECT:		POT-HYI	FBOS/	BI AIR SI	IP ONE (I	MURRAY	PACIFIC	3)		USACE	GRAYS H.	O&M
DMMU ID:	C20	C21	C22	C23	C24	C25	C27	C36	C37	C5	C10	C11
Testing Rank:	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н	Н
METALS & ORGANOMETALS												
Antimony												
Arsenic		S										
Cadmium												
Copper												
Lead												
Mercury												
Nickel												
Silver		-										
Zinc	OD	SB (G)		SB (G)	00 (0)				SB (G)			
TBT ion (porewater) LPAH	SB	SB (G)		SB (G)	SB (G)				SB (G)			
Naphthalene												
Acenaphthene												
Acenaphthylene												
Fluorene												
Phenanthrene												
Anthracene												
2-Methylnaphthalene												
Total LPAHs												
HPAH												
Fluoranthene					S	SB	S					
Pyrene					S	S						
Benzo(a)anthracene					S	S						
Benzofluoranthenes (b+k)					S	S						
Chrysene	S				S	S	S					
Benzo(a)pyrene				ļ		ļ						
Indeno(1,2,3-c,d)pyrene												
Dibenzo(a,h)anthracene												
Benzo(g,h,l)perylene	-	1		-					 			
Total HPAHs					S	S						
CHLORINATED HYDROCARBONS			-	 		 			 		-	
1,2,4-Trichlorobenzene												
1,2-Dichlorobenzene												
1,3-Dichlorobenzene 1,4-Dichlorobenzene												
Hexaclorobenzene (HCB)		S (U)		S (U)							S (U)	
PHTHALATES		3 (0)		3 (0)							3 (0)	
Bis(2-ethylheyesyl)phthalate												
Butylbenzylphthalate												
Di-n-butylphthalate												
Di-n-octylphthalate												
Diethylphthalate												
Dimethylphthalate												
PHENOLS												
2-Methylphenol												
4-Methylphenol												
2,4-Dimethylphenol		S (U)		S(U)							S(U)	
Pentachlorophenol												
Phenol												
MISCELANEOUS EXTRACTABLES												
Benzyl alcohol												
Benzoic acid	-	1		-		-			 			
Dibenzofuran	l	0.410		0.410		-			 		C/LIV	
Hexachlorobutadiene		S (U)	-	S (U)		 			 		S(U)	
Hexachloroethane N-Nitrosodiphenylamine		S (U)		S (U)		-			-			
VOLATILE ORGANICS	-	3 (0)		3 (0)		 			 			
Ethylbenzene	-											
Tetrachloroethene		†		1		1						
Total Zylene (sum of o,m,p)												
Trichloroethane												
PESTICIDES AND PCBs												
Total DDT	S	S	S	S (UE)	S (UE)	S (E)	S	S	S			
Aldrin				,,	,/	_/						
alpha-Chlordane												
Dieldrin	S (Y)		S (Y)		S	S (Y)	S					
Heptachlor												
gamma-BHC (Lindane)	S (Y)		S (E)									
Total PCBs	S	S		S	S	S	S	S	S			
BIOASSAYS												
Amphipod						1H			2H			
Sediment Larval (Bivalve/Echinoderm)	1H	1H	1H	1H	1H	2H	2H	1H				
Neanthes Growth												
Bioassay: (Pass/Fail)	FAIL	FAIL	FAIL	FAIL	FAIL	FAIL	PASS	FAIL	PASS	PASS	NA	PASS
BTs eyesceeded:	yes	yes	no	yes	yes	yes	no	no	yes	no	no	no
Bioaccumulation test conducted:	no	no		no	no	no			no			
Bioaccumulation (Pass/Fail):												
ML Rule exceeded:	no	no	no	no	no	no	no	no	no	no	no	no
OVERALL PASS/FAIL:	FAIL	FAIL	FAIL	FAIL	FAIL	FAIL	PASS	FAIL	FAIL(C)	PASS	PASS (BPJ)	PASS
HIGHEST RANKING:	H*	H*	H*	H*	H*	H*	LM	H*	H*	L	LM	L